

ERRATUM Open Access



Erratum: Uterine NDRG2 expression is increased at implantation sites during early pregnancy in mice, and its down-regulation inhibits decidualization of mouse endometrial stromal cells

Yan Gu¹, Xuan Zhang^{2*}, Qian Yang¹, Jian-mei Wang³, Ya-ping He², Zhao-gui Sun², Hui-qin Zhang¹ and Jian Wang^{2*}

After publication of this article [1], the authors noticed an error in Fig. 5: in panel c, the bands produced by NDRG2 and beta-actin were incorrectly labelled. The original version of this article has been updated to correct this. The correct figure is shown below:

Author details

¹Shanghai Medical School, Fudan University, Shanghai, China. ²NPFPC Key Laboratory of Contraceptive Drugs & Devices, Shanghai Institute of Planned Parenthood Research, Shanghai, China. ³The Second Hospital of Tianjin Medical University, Tianjin, China.

Received: 3 August 2015 Accepted: 3 August 2015 Published online: 29 August 2015

Reference

 Gu Y, Zhang X, Yang Q, Wang JM, He YP, Sun ZG, et al. Uterine NDRG2 expression is increased at implantation sites during early pregnancy in mice, and its down-regulation inhibits decidualization of mouse endometrial stromal cells. Reprod Biol Endocrinol. 2015;13:49.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit



²NPFPC Key Laboratory of Contraceptive Drugs & Devices, Shanghai Institute of Planned Parenthood Research, Shanghai, China



^{*} Correspondence: zhangxuancw@outlook.com; wangjiansippr@126.com The online version of the original article can be found under doi:10.1186/s12958-015-0047-7.




Fig. 5 In vitro decidualization of mouse primary ESCs. ESCs isolated from the uteri of day 4 pregnant mice were cultured in the presence of P4 and E2. **a** Immunocytochemical detection of vimentin and cytokeratin in cultured ESCs. **b** Quantitative PCR analysis of DTPRP mRNA expression in ESCs cultured for up to 72 h. **c** Western blot analysis of NDRG2 protein expression in ESCs cultured for up to 72 h. Densitometric analyses of NDRG2 at each time point compared with 0 h is shown. **d** Quantitative PCR analysis of NDRG2 mRNA expression in cultured ESCs. The relative fold induction of NDRG2 mRNA expression at each time point compared with its expression in the 0 h sample is shown. The values represent the mean ± SEM, as determined from three separate experiments. *, significantly different (P < 0.05)