

RESEARCH

Open Access



Risk factors for missed abortion: retrospective analysis of a single institution's experience

Wei-Zhen Jiang¹, Xi-Lin Yang² and Jian-Ru Luo^{1*}

Abstract

Objective: To explore the risk factors including the difference between mean gestational sac diameter and crown-rump length for missed abortion.

Methods: Hospitalized patients with missed abortion and patients with continuing pregnancy to the second trimester from Chengdu Women's and Children's Central Hospital from June 2018 to June 2021 were retrospectively analyzed. The best cut-off value for age and difference between mean gestational sac diameter and crown-rump length (mGSD-CRL) were obtained by x-tile software. Univariate and multivariate logistic regression analysis were adopted to identify the possible risk factors for missed abortion.

Results: Age, gravidity, parity, history of cesarean section, history of recurrent abortion (≥ 3 spontaneous abortions), history of ectopic pregnancy and overweight or obesity ($\text{BMI} > 24 \text{ kg/m}^2$) were related to missed abortion in univariate analysis. However, only age (≥ 30 vs < 30 years: $\text{OR} = 1.683$, $95\% \text{CI} = 1.017-2.785$, $P = 0.043$, power = 54.4%), BMI (> 24 vs $\leq 24 \text{ kg/m}^2$: $\text{OR} = 2.073$, $95\% \text{CI} = 1.056-4.068$, $P = 0.034$, power = 81.3%) and mGSD-CRL (> 20.0 vs $\leq 11.7 \text{ mm}$: $\text{OR} = 2.960$, $95\% \text{CI} = 1.397-6.273$, $P = 0.005$, power = 98.9%; $11.7 < \text{mGSD-CRL} \leq 20.0$ vs $> 20.0 \text{ mm}$: $\text{OR} = 0.341$, $95\% \text{CI} = 0.172-0.676$, $P = 0.002$, power = 84.8%) were identified as independent risk factors for missed abortion in multivariate analysis.

Conclusion: Patients with age ≥ 30 years, $\text{BMI} > 24 \text{ kg/m}^2$ or $\text{mGSD-CRL} > 20 \text{ mm}$ had increasing risk for missed abortion, who should be more closely monitored and facilitated with necessary interventions at first trimester or even before conception to reduce the occurrence of missed abortion to have better clinical outcomes.

Keywords: First trimester pregnancy, Missed abortion, Gestational sac, Crown-rump length, mGSD-CRL

Background

Missed abortion was a special type of spontaneous abortion that the embryo or fetus has already died but remained in the uterus for days or weeks and with a closed cervical ostium [1]. Patients might present with or

without subtle clinical symptoms such as vaginal bleeding or abdominal pain. Missed abortion, occurring in approximately 8–20% of clinically confirmed intrauterine pregnancies [2], was often confirmed using ultrasonography.

Missed abortion was undoubtedly a huge physical and psychological setback for women with fertility requirements. Therefore, early identification of women at high risk of missed abortion was pivotal, which might aid in providing possible theoretical basis for implementing clinical measures to prevent missed abortion. Previous studies have revealed that Human Chorionic Gonadotropin (HCG), Estradiol (E2),

*Correspondence: 1032261258@qq.com

¹ Department of Gynaecology and Obstetrics, Chengdu Women's and Children's Central Hospital, School of Medicine, University of Electronic Science and Technology of China, No.1617 of Riyue Avenue, Qingyang District, Chengdu 611731, China
Full list of author information is available at the end of the article



progesterone, gestational sac diameter (GSD), Crown-Rump Length (CRL), fetal heart rate and yolk sac diameter might be predictive for early pregnancy loss [3–5]. In addition, the predictive value of mGSD-CRL for early pregnancy outcome in in vitro fertilization (IVF) treatment has been established [6]. However, most of the current studies have performed univariate analysis to identify the risk factors for early pregnancy loss [3–6].

Therefore, we conducted this study to more comprehensively explore the possible high risk factors relating to developing of missed abortion using multivariate logistic regression analysis, hopefully it could be of great help to identification and intervention.

Materials and methods

Data sources

We reviewed patients from Chengdu Women's and Children's Central Hospital from June 2018 to June 2021. Inclusion criteria of missed abortion group were listed as follows: (1) Not more than 12 weeks gestation; (2) Crown-rump length \geq seven mm without heartbeat or (3) mean sac diameter \geq 25 mm without embryo or (4) absence of embryo with heartbeat \geq two weeks after a scan that showed a gestational sac without a yolk sac or (5) absence of embryo with heartbeat \geq 11 days after a scan that showed a gestational sac with a yolk sac [7]. Exclusion criteria of missed abortion group were listed as follows: (1) Incomplete information; (2) multiple pregnancy. Patients with the following inclusion and exclusion criteria were enrolled as control group: (1) Patients continued pregnancy to the second trimester were included; (2) Incomplete information and multiple pregnancy were excluded. After excluding patients with incomplete information, 307 patients were finally included with 160 patients having missed abortion and 147 with continuing pregnancy to second trimester. Due to the retrospective nature of the study, informed consent was waived, but this study was granted by the ethics committee of Chengdu Women's and Children's Central Hospital and the ethics approval number was B2021(26).

Collection of data

Patients' information regarding age, gravidity, parity, history of vaginal delivery, history of cesarean delivery, history of recurrent abortion (\geq 3 spontaneous abortions), history of induced abortion, history of medication abortion, history of midtrimester induction, history of ectopic pregnancy, history of smoking, history of alcohol consumption, history of other uterine operations, mode of conception, BMI, mGSD-CRL not more than 12 weeks with live embryo were collected.

Statistical analysis

Categorical variables were described as percentages or frequencies and compared using Pearson χ^2 test; continuous variables were described as medians with interquartile range (IQR) and compared with *t* test. We identified the cut-off value for age and mGSD-CRL via X-tile software (version 3.6.1; Yale University, New Haven, CT, USA) once maximal chi-square value reached, which was considered to represent the greatest difference in outcomes prediction among the subgroups [8].

Logistic regression was used to determine independent risk factors for missed abortion. Statistically significant variables from univariate logistic regression analysis ($P < 0.1$) were included in the multivariate analysis. Pearson χ^2 test, *t* test and logistic regression were performed using SPSS (version 25.0, SPSS, Chicago, IL, USA), X-tile software was used to calculate cut-off value. G*Power Analysis program (version 3.1, The G*Power Team, Belgium) was used for power calculation. A two-tailed $P < 0.05$ was recognized as statistically significant.

Results

Study cohort

A total of 307 patients were finally included in the study with 160 cases having missed abortion and 147 with continuing pregnancy to second trimester (Supplementary Fig. 1). The characteristics were listed in Supplementary Table 1. As a result, 30 years old was the cut-off value for age via X-tile software. Therefore, age was split as age \geq 30 years and age $<$ 30 years. Similarly, mGSD-CRL was divided into three subgroups: GSD-CR $<$ 11.7 mm, $11.7 \text{ mm} \leq \text{mGSD-CRL} \leq 20.0 \text{ mm}$, GSD-CR $>$ 20.0 mm (Fig. 1). Nearly half of the patients were over 30 years old (49.2%). 38.4% of the patients were having first pregnancy, and the majority of the patients had never delivered (71.0%). 11.1% of the patients had a history of vaginal delivery, however, 18.2% of the patients had a history of cesarean section. Of note, 16% of the patients had a BMI $>$ 24 kg/m², 29.6% of the patients had a mGSD-CRL $<$ 11.7 mm and 18.9% had a mGSD-CRL $>$ 20 mm. Moreover, 2.6% of the patients suffering from recurrent abortion and 4.2% had a history of ectopic pregnancy. Besides, 39.1% of the patients had a history of curettage. In total, 52.1% of the patients developed missed abortion (Table 1).

Risk factors for missed abortion

In the univariate logistic regression analysis, Age, gravidity, parity, history of cesarean section, history of recurrent abortion, history of ectopic pregnancy, overweight or obesity (BMI $>$ 24 kg/m²) and mGSD-CRL




Fig. 1 mGSD-CRL at diagnosis stratification by X-tile software

were significantly related to increased risk factors for missed abortion. Furthermore, risk factors identified in the univariate logistic regression analysis were included in the multivariate analysis, which revealed that Age (≥ 30 vs < 30 years: OR = 1.683, 95%CI = 1.017–2.785, $P = 0.043$, power = 54.4%), BMI (> 24 vs ≤ 24 kg/m²: OR = 2.073, 95%CI = 1.056–4.068, $P = 0.034$, power = 81.3%), mGSD-CRL (> 20.0 vs ≤ 11.7 mm: OR = 2.960, 95% CI = 1.397–6.273, $P = 0.005$, power = 98.9%; $11.7 < \text{mGSD-CRL} \leq 20.0$ vs > 20.0 mm: OR = 0.341, 95%CI = 0.172–0.676, $P = 0.002$, power = 84.8%) were independent risk factors for missed abortion (Table 2).

Discussion

Missed abortion, normally presenting without symptoms of threatened abortion such as abdominal pain and vaginal bleeding, was a kind of spontaneous abortion, which were frequently diagnosed using ultrasonography. In this study, we retrospectively analyzed the data of 160 missed abortion patients and 147 pregnant women who didn't have abortion in the first trimester in order to fully establish the possible risk factors for missed abortion, and

provide evidence for early identification and intervention for patients with high risk of missed abortion.

In previous studies, it was believed that advanced age was a high risk factor for missed abortion, which might result from the decline of ovarian function and corpus luteum function as age accrued [1, 9]. However, previous study also showed that advanced age was not a high risk factor for spontaneous abortion [10], in which age was divided into advanced age group (> 35 years) and non-advanced age group (≤ 35 years old). Therefore, we hypothesized that there might be a more meaningful cutoff value other than 35 years old to divide the age into two subgroups. As a result, 30 years old, calculated via x-tile, showed significant value in the final multivariate logistic analysis (OR = 1.683, 95%CI = 1.017–2.785, $P = 0.043$). As controversial regarding age existed in previous studies, our result showing that age > 30 was an independent risk factor for missed abortion seemed solid. And the dropping from 35 to 30 in terms of cut-off value for age might be related to factors like increasing pressure, unhealthy living habits and environmental pollution resulting from social developing [2, 11]. Although

Table 1 The characteristics of risk factors

Variables	Count(%)	Variables	Count(%)
Age(years)		History of medication abortion	
30	156(50.8%)	Yes	6(2.0%)
≥ 30	151(49.2%)		
Gravidity(times)		History of midtrimester induction	
1	118(38.4%)	Yes	5(1.6%)
2	79(25.7%)		
3	53(17.3%)	History of ectopic pregnancy	
4	36(11.7%)	Yes	13(4.2%)
5	13(4.2%)		
6	7(2.3%)	Smoking	
7	1(0.3%)	Yes	1(0.1%)
Parity(times)		Alcohol consumption	
0	218(71.0%)	Yes	1(0.1%)
1	83(27.0%)		
2	6(2.0%)	Other uterine operations	
History of vaginal delivery		Yes	7(2.3%)
Yes	34(11.1%)		
History of cesarean delivery		IVF	
Yes	56(18.2%)	Yes	9(2.9%)
History of recurrent abortion		BMI(Kg/m²)	
Yes	8(2.6%)	≤ 24	258(84%)
		> 24	49(16%)
History of induced abortion		mGSD-CRL(mm)	
Yes	120(39.1%)	< 11.7	91(29.6%)
		11.7 ≤ mGSD-CRL ≤ 20.0	158(51.5%)
		> 20.0	58(18.9%)

the cut-off value in our study were not consistent with previous ones, the consensus on older age was a high risk factor for missed abortion was basically reached.

A meta-analysis including 16 studies demonstrated that BMI > 25 kg/m² was a high risk factor for abortion [12], which reported that the missed abortion rate of overweight or obese women was as high as 25–37% [13]. The participants from our study were childbearing age women from China, so the definition of overweight or obese as BMI > 24 kg/m² was used for grouping though the World Health Organization(WHO) defined overweight or obesity as BMI > 25 kg/m² [14]. And the result showed that patients with BMI > 24 kg/m² were more likely to have missed abortion than BMI ≤ 24 kg/m² (OR = 2.073, 95% CI = 1.056–4.068, P = 0.034), which was consistent with previous studies [11, 12]. Therefore, weight control before pregnancy was usually recommended.

Although the effect of mGSD and CRL on missed abortion had been reported [3, 4, 15–18], there was few studies working on the relationship between mGSD-CRL and missed abortion. Bromley et al. firstly proposed the concept of "small gestational sac" [19]. And their work

revealed that mGSD-CRL < 5 mm in the first trimester was a high risk factor for missed abortion. However, the extremely limited number of included patients in their study might impede the generalization of the conclusion. Similarly, the research from Kapfhamer et al. also showed that mGSD-CRL < 5 mm was a high risk factor for early pregnancy loss, and further demonstrated that mGSD-CRL > 10 mm was a protective factor for early pregnancy loss [6]. However, Zhao et al. believed that "large gestational sac"(mGSD-CRL ≥ 18 mm) was related to increasing risk for spontaneous abortion [20]. Therefore, we used x-tile to find the two optimal cut-off values for mGSD-CRL, which showed that patients with mGSD-CRL > 20 mm was more more likely to have missed abortion than patients with mGSD-CRL ≤ 20 mm. And there was no statistical difference between mGSD-CRL < 11.7 mm group and 11.7 ≤ mGSD-CRL ≤ 20.0 mm group. In summary, we were inclined to believe that increasing mGSD-CRL was associated with increasing risk of missed abortion, which should be further validated in the future due to the differences in sample size from previous studies [6, 19, 20].

Table 2 Logistic regression analysis

Variables	OR(95%CI)	P-value	OR(95%CI)	P-value
	univariate logistic regression analysis		multivariate logistic regression analysis	
Age(years)				
30	Reference		Reference	
≥ 30	2.018(1.280–3.181)	0.002	1.683(1.017–2.785)	0.043
Gravidity(times)				
1	Reference			
2	1.009(0.570–1.786)	0.975		
3	1.384(0.722–2.653)	0.328		
4	2.291(1.048–5.006)	0.038		
5	2.577(0.752–8.836)	0.132		
6	2.864(0.534–15.353)	0.219		
7	1,850,453,001.811(-)	1.000		
Parity(times)				
0	Reference			
1	2.079(1.231–3.510)	0.006		
2	1.117(0.220–5.654)	0.894		
History of cesarean delivery				
No	Reference			
Yes	2.232(1.209–4.121)	0.010		
Recurrent abortion				
No	Reference			
Yes	6.680(0.812–54.960)	0.077		
History of ectopic pregnancy				
No	Reference			
Yes	3.200(0.863–11.863)	0.082		
BMI(Kg/m²)				
≤ 24	Reference		Reference	
24	1.912(1.011–3.614)	0.046	2.073(1.056–4.068)	0.034
mGSD-CRL(mm)				
11.7	Reference		Reference	
11.7 ≤ mGSD-CRL ≤ 20.0	0.941(0.561–1.577)	0.818	1.009(0.579–1.758)	0.976
20.0	2.804(1.382–5.689)	0.004	2.960(1.397–6.273)	0.005
20.0			Reference	
11.7			0.338(0.159–0.716)	0.005
11.7- 20.0			0.341(0.172–0.676)	0.002

Age, gravidity, parity, history of cesarean section, history of recurrent abortion, history of ectopic pregnancy, BMI and mGSD-CRL were identified in the univariate analysis. However, only age, BMI and mGSD-CRL were still meaningful in multivariate analysis. What was inconsistent with previous studies in our study was that recurrent abortion was not a high risk factor for missed abortion [21], which might result from the low incidence of recurrent abortion (missed abortion group vs non-missed abortion group: 7 vs 1) in our study.

One major strength of this study was that stratifying age by x-tile rather than 35 years were firstly recognized

for high risk of missed abortion. Other strengths included that mGSD-CRL were analyzed instead of mGSD or CRL independently. On the contrary, This study was inevitably limited by the retrospective nature. In addition, the pathogenic factors for missed abortion was complicated, and some possible high risk factors like immunological or genetic factors could not be obtained.

It is well known that missed abortion is a special type of spontaneous abortion and the ultimate outcome is embryonic arrest. The current knowledge of the missed abortion mostly relates to prevention and treatment, but the classification and severity have not been covered yet

according to existing literature and guidelines. The purpose of this paper is to explore the high-risk factors of missed abortion, therefore treatment was barely involved, and we will do more research on the treatment of missed abortion in future work. Overall, We hope that the present study could aid in abortion prediction and treatment decision-making for clinicians.

Conclusions

This study demonstrated that age ≥ 30 years old, BMI > 24 kg/m² and mGSD-CRL > 20 mm were independent risk factors for missed abortion. This study provided a theoretical basis for clinicians to deliver prompt interventions in childbearing age women during the first trimester or even before pregnancy, so as to reduce the incidence of missed abortion.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12958-022-00987-2>.

Additional file 1: Supplementary Figure 1. Flow chart depicting for inclusion of study subjects. **Supplementary Table 1.** Clinical characteristics of participants.

Acknowledgements

The authors wish to thank the participants, Chengdu Women's and Children's Central Hospital hospital staff, and whoever contributed to this study.

Authors' contributions

Conception/design: Wei-Zhen Jiang. Provision of study material or patients: Wei-Zhen Jiang. Collection and/or assembly of data: Wei-Zhen Jiang, Xi-Lin Yang. Data analysis and interpretation: Wei-Zhen Jiang, Xi-Lin Yang. Manuscript writing: Wei-Zhen Jiang. Manuscript revision: Jian-Ru Luo. Final approval of manuscript: All authors.

Funding

Not applicable.

Availability of data and material

All data that support the findings of this study were available from the corresponding author via E-mail due to appropriate request.

Declarations

Ethics approval and consent to participate

Due to the retrospective nature of the study, informed consent was waived, but this study was granted by the ethics committee of Chengdu Women's and Children's Central Hospital and the ethics approval number was B2021(26).

Consent for publication

Written informed consent for publication was obtained from all participants.

Competing interests

The authors have declared that there is on potential conflicts of interest.

Author details

¹Department of Gynaecology and Obstetrics, Chengdu Women's and Children's Central Hospital, School of Medicine, University of Electronic Science and Technology of China, No.1617 of Riyue Avenue, Qingyang District, Chengdu 611731, China. ²Department of Radiation Oncology, Chengdu Women's and Children's Central Hospital, School of Medicine, University of Electronic Science and Technology of China, Chengdu 611731, China.

Received: 9 January 2022 Accepted: 3 August 2022

Published online: 09 August 2022

References

- Cunningham FG, Leveno KJ, Bloom SL, Dashe JS, Hoffman BL, Casey BM, et al. *Williams Obstetrics*. 25nd ed. New York: McGraw-Hill; 2018.
- Fang J, Xie B, Chen B, et al. Biochemical clinical factors associated with missed abortion independent of maternal age: A retrospective study of 795 cases with missed abortion and 694 cases with normal pregnancy. *Medicine (Baltimore)*. 2018;97(50):e13573.
- Altay MM, Yaz H, Haberal A. The assessment of the gestational sac diameter, crown-rump length, progesterone and fetal heart rate measurements at the 10th gestational week to predict the spontaneous abortion risk. *J Obstet Gynaecol Res*. 2009;35(2):287–92.
- Papaioannou GI, Syngelaki A, Maiz N, et al. Ultrasonographic prediction of early miscarriage. *Hum Reprod*. 2011;26(7):1685–92.
- Puget C, Joueidi Y, Bauville E, et al. Serial hCG and progesterone levels to predict early pregnancy outcomes in pregnancies of uncertain viability: A prospective study. *Eur J Obstet Gynecol Reprod Biol*. 2018;220:100–5.
- Kapfhamer JD, Palaniappan S, Summers K, et al. *Fertil Steril*. 2018;109(1):130–6.
- Doubilet PM, Benson CB, Bourne T, et al. Diagnostic criteria for nonviable pregnancy early in the first trimester. *N Engl J Med*. 2013;369(15):1443–51.
- Zhuang W, Chen J, Li Y, et al. Valuation of lymph node dissection in localized high-risk renal cell cancer using X-tile software. *Int Urol Nephrol*. 2020;52(2):253–62.
- Guifang G, Caixin Y, Yanqing H, et al. A survey of influencing factors of missed abortion during the two-child peak period. *J Obstet Gynaecol*. 2020;undefined:1–4.
- Yavuz P, Taze M, Salihoglu O. The effect of adolescent and advanced-age pregnancies on maternal and early neonatal clinical data. *J Matern Fetal Neonatal Med*. 2021;undefined:1–7.
- Wu J, Hou H, Ritz B, et al. Exposure to polycyclic aromatic hydrocarbons and missed abortion in early pregnancy in a Chinese population. *Sci Total Environ*. 2010;408(11):2312–8.
- Metwally M, Ong KJ, Ledger WL, et al. Does high body mass index increase the risk of miscarriage after spontaneous and assisted conception? A meta-analysis of the evidence. *Fertil Steril*. 2008;90(3):714–26.
- Hamilton-Fairley D, Kiddy D, Watson H, et al. Association of moderate obesity with a poor pregnancy outcome in women with polycystic ovary syndrome treated with low dose gonadotrophin. *Br J Obstet Gynaecol*. 1992;99(2):128–31.
- WHO. Obesity: preventing and managing the global epidemic. *World Health Organ Tech Rep Ser*. 2000;894:1–253.
- Pexsters A, Luts J, Van Schoubroeck D, et al. Clinical implications of intra- and interobserver reproducibility of transvaginal sonographic measurement of gestational sac and crown-rump length at 6–9 weeks' gestation. *Ultrasound Obstet Gynecol*. 2011;38(5):510–5.
- Preisler J, Kopeika J, Ismail L, et al. Defining safe criteria to diagnose miscarriage: prospective observational multicentre study. *BMJ*. 2015;351:4579.
- Ouyang Y, Qin J, Lin G, et al. Reference intervals of gestational sac, yolk sac, embryonic length, embryonic heart rate at 6–10 weeks after in vitro fertilization-embryo transfer. *BMC Pregnancy Childbirth*. 2020;20(1):533.
- Papaioannou GI, Syngelaki A, Poon LCY, et al. Normal ranges of embryonic length, embryonic heart rate, gestational sac diameter and yolk sac diameter at 6–10 weeks. *Fetal Diagn Ther*. 2010;28(4):207–19.
- Bromley B, Harlow BL, Laboda LA, et al. Small sac size in the first trimester: a predictor of poor fetal outcome. *Radiology*. 1991;178(2):375–7.
- Xuan Z, Liu Lu, Haiqian Li. Clinical Analysis of the Impact of Corpus Luteum, Gestational Sac and Embryo in Early Pregnancy on the Pregnant Outcomes. *J Pract Obstet Gynecol*. 2013;08:595–8.
- Li H, Qin S, Xiao F, et al. Predicting first-trimester outcome of embryos with cardiac activity in women with recurrent spontaneous abortion. *J Int Med Res*. 2020;48(6):300060520911829.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.