

Review

Open Access

Aromatase expression and role of estrogens in male gonad : a review

Serge Carreau*¹, Sophie Lambard¹, Christelle Delalande¹, Isabelle Denis-Galeraud², Barbara Bilinska³ and Sonia Bourguiba¹

Address: ¹Department of Biochemistry, University, Esplanade de la Paix, Caen, France, ²Department of genetic & reproduction, CHU Clémenceau, Caen, France and ³Department of Zoology, University Jagiellonian, Cracow, Poland

Email: Serge Carreau* - carreau@ibba.unicaen.fr; Sophie Lambard - solambard@yahoo.fr; Christelle Delalande - delalande@ibba.unicaen.fr; Isabelle Denis-Galeraud - denis-i@chu-caen.fr; Barbara Bilinska - bbili@z.uk.iz.uj.edu.pl; Sonia Bourguiba - sbourguiba@yahoo.fr

* Corresponding author

Published: 11 April 2003

Received: 20 March 2003

Reproductive Biology and Endocrinology 2003, **1**:35

Accepted: 11 April 2003

This article is available from: <http://www.RBEj.com/content/1/1/35>

© 2003 Carreau et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

The ability of the testis to convert irreversibly androgens into estrogens is related to the presence of a microsomal enzymatic complex named aromatase, which is composed of a specific glycoprotein, the cytochrome P450 aromatase (P450arom) and an ubiquitous reductase. The aromatase gene is unique in humans and contained 18 exons, 9 of them being translated. In the rat testis we have immunolocalized the P450arom not only in Leydig cells but also in germ cells and especially in elongated spermatids. Related to the stage of germ cell maturation, we have shown that the level of P450arom mRNA transcripts decreases, it is much more abundant in pachytene spermatocytes and round spermatids than in mature germ cells whereas the aromatase activity is 2–4 fold greater in spermatozoa when compared to the younger germ cells. Using a highly specific quantitative competitive RT-PCR method we have evidenced that several factors direct the expression of the aromatase gene in Leydig cells, Sertoli cells, pachytene spermatocytes and round spermatids, and it is obvious that promoter PII is the main one but other promoters could be concerned.

In the bank-vole testis we have observed a positive correlation between a fully developed spermatogenesis and a strong immunoreactivity for both P450arom and estrogen receptor β not only in Sertoli cells but also in pachytene spermatocytes and round spermatids. Our recent data obtained from ejaculated human spermatozoa demonstrate the presence of aromatase both in terms of mRNA and protein, and in addition, we suggest that aromatase could be involved in the acquisition of sperm motility. Indeed in men the congenital aromatase deficiency is associated with severe bone maturation problems and sterility. Together with the widespread distribution of estrogen receptors in testicular cells these data clearly show that estrogens play a physiological role in the regulation of spermatogenesis in mammals.

Introduction

The mammalian testis is a complex organ characterised by two main functions : synthesis of steroid hormones and production of spermatozoa. It is well known that normal

testicular development and maintenance of spermatogenesis are controlled by gonadotrophins and testosterone whose effects are modulated by locally-produced factors, and among them estrogens are obviously concerned [1].

Within the reproductive tract of the male the levels of estrogens are far higher than in the general blood compartment [2], therefore favoring a testicular source of estrogens [3].

The role of estrogens in the development and physiology of male reproductive tract of mammals is still a matter of debates even though there is a growing body of evidence suggesting that estrogens are playing a role via their specific receptors (ER α and ER β) which are distributed all along the genital tract [4–6].

Aromatase expression in male germ cells

The cytochrome P450 aromatase (P450arom) is involved in the irreversible transformation of androgens into estrogens and is present in the endoplasmic reticulum of numerous vertebrate tissues. The P450arom is a microsomal enzymatic complex composed of two proteins: a ubiquitous NADPH-cytochrome P450 reductase and a cytochrome P450 aromatase, which contains the heme and the steroid binding pocket. In humans the P450arom is the product of a single gene called *Cyp19*, which belongs to the cytochrome P450 gene family, containing more than 500 members [7]. The *Cyp19* gene lies on more than 120 kb length with a coding region of 9 exons plus 9 untranslated exons I. The *Cyp19* gene expression is regulated by tissue-specific promoters producing alternate 5'-untranslated exons I that are then spliced onto a common 3'-splice acceptor site in the exon II upstream of the translation start site [8].

In the mammalian testis it's well known that aromatase is mainly localized in Leydig cells (for review [1]). In rodents, past efforts to determine the source of testicular estrogens have been a considerable subject of interest and led to the assertion that Leydig cells synthesize estrogen in adults, whereas Sertoli cells are the major source in immature animals. However it has been shown that germ cells from adult male rat represent a new source of estrogens [4]. Indeed, the amount of P450arom transcripts is higher in pachytene spermatocytes than in round spermatids and obviously than in testicular spermatozoa. Conversely the aromatase activity is more intense in haploid germ cells than in younger germ cells which was also confirmed by the immunolocalisation of the protein on testicular cells mainly elongated spermatids [9]. These observations have been corroborated by Janulis et al [10] who showed that the aromatase activity is localized in the cytoplasmic droplet and decreased according to the epididymal region, being higher in caput than in cauda. Moreover it has been reported the existence of P450arom transcripts in the epithelial cells of the rat epididymis [11].

In mouse [12] as well as in bank vole [13], the aromatase is present not only in Leydig cells and Sertoli cells but also

in germ cells. In the bank vole, a seasonal breeder, the P450arom is much more expressed (in terms of specific transcripts, activity and immunolocalisation) in pachytene spermatocytes and spermatids of animals bred in long day-light cycle than in animals in winter rest [13,14]. Indeed there is a synchronisation between the recrudescence of spermatogenesis and the levels of expression of aromatase and estrogen receptors beta. Recent observations made in our group showed that in seminiferous tubules of the stallion there was a positive immunoreactive signal for aromatase not only in Leydig cells but also in cytoplasm surrounding germ cells therefore suggesting strongly the presence of aromatase in Sertoli cells [15].

In the *Rhesus* monkey, it has been reported that testis and to a lesser extent epididymidis contained P450arom transcripts [16] and moreover, in the epididymal regions a discrepancy was observed between the amount of transcripts and the aromatase activity which was found more active in caput than in cauda although the P450arom mRNA levels were reversed. In humans the main source of estrogens is in Leydig cells; nevertheless the Sertoli cells are able to synthesize estradiol *in vitro* [17]. It has been also claimed that estrogens are present in spermatozoa [1,18]. From these observations we have examined the ability of human ejaculated spermatozoa to transform androgens into estrogens. When sperm RNA was used as template in RT-PCR we have shown the presence of P450arom transcripts; the sequences alignment from these PCR products and granulosa cells with published human P450arom gene were identical. Using Western blots and a specific monoclonal antibody against aromatase [19] we have evidenced the presence of aromatase in these sperm cells which was obviously more abundant in spermatozoa containing cytoplasmic droplets. These observations are in fitting with other recent data [20], and in addition we have demonstrated that the amount of P450arom transcripts was 30% lower in immotile than in motile spermatozoa [21].

In order to bring insights onto the role of estrogens in seminiferous tubules, it is first necessary to analyse the regulation of the *Cyp19* gene transcription. Thus we have evidenced using RACE-PCR that promoter II directs the expression of aromatase gene whatever the testicular cell type studied in the rat [22]. Germ cells are potent targets for growth factors and cytokines; indeed, pachytene spermatocytes (PS) and round spermatids (RS) produce TGF β and TNF α ; but only TGF β receptors are present in germ cells [23]. Accordingly in highly purified germ cells from adult rat (pachytene spermatocytes and round spermatids) we have demonstrated that TGF β inhibits the expression of *Cyp19* in both germ cell fractions although TNF α exerts a stimulatory role in pachytene spermatocytes only; these data have been confirmed by the estradiol outputs

Table 1: Aromatase deficiencies in men

Authors	Affected Exon	Mutation
Morishima et al., 1995*	IX	Arg375Cys
Carani et al., 1997	IX	Arg365Gln
Deladoey et al., 1999*	V	Leu157XΔC-stop
Maffei et al., 2001*	V	Nucleotide 628 G to A Insertion of 10aa and stop
Kottler et al., 2002	IX	Insertion of T position 1058 and stop
Herrmann et al., 2002	V	C to A substitution position 3 and stop

* see Carani et al [44] for details




Figure 1

Aromatase and estrogen receptors (ER) in adult male rat gonad. PS: pachytene spermatocytes, RS: round spermatids, Spz: spermatozoa. Aromatase has been demonstrated in terms of mRNA (RT-PCR), protein (Western blots) and enzyme activity (measurements of estradiol output in culture media) in the various testicular cells. ER : estrogen receptors localisation.




Figure 2
Human spermatozoa : source and targets of estrogens. It has been shown that human ejaculated spermatozoa contain aromatase as revealed by the mRNA, protein on Western blots and the presence of endogenous estradiol. As far as estrogen receptors are concerned mainly ER alpha are present.

measured in germ cell culture media [23]. It is noteworthy that the effect of TNF α is amplified in presence of dexamethasone which is may be supported by the presence of an other promoter like promoter I.4 as shown in adipose tissue [24]. It is also obvious that other modulators direct the expression of the aromatase gene since in Leydig cells we have shown that the amount of P450arom transcripts are increased in presence of seminiferous conditioned medium [25]. Our data are likely in agreement with a recent work showing that the aromatase expression is controlled by three different promoters in mice that is to say that in the male gonad the *Cyp19* gene expression is directed by a testis specific promoter [26].

Estrogens and testicular functions

In order to exert a biological role, testicular estrogens should interact with estrogen receptors (ER) which in turn modulate the transcription of specific genes. Therefore considering the presence of at least two ER (ER α and ER β) in most of the testicular cells and the other parts of the genital tract, the physiological role of estrogens in male reproduction is now extensively revisited [4–6,27,28]. Spermatogenesis in rodents is in part under estrogens control, namely the stem germ cell number and the spermatid maturation [29,30]. In the bank the expression of androgen receptors, P450arom and ERs (α and β) in testicular cells is related to the length of the photoperiod. More precisely P450arom and ER β are much more expressed in testes (especially in spermatocytes and elongated spermatids) of long photoperiod-reared animals in which

spermatogenesis is fully developed when compared to short-day length bred animals with regressed testes [13,31]. These observations are in fitting with previous reports [32,33] and the recent observations of Pak et al [34] demonstrating an improvement of the recrudescence of spermatogenesis in estradiol-treated hamster which have been confirmed by Bilinska et al [35] in bank vole.

Therefore estradiol is now considered as a survival factor for germ cells [36] that is consistent with data obtained in male monkeys treated with aromatase inhibitor in which a blockage of spermatid maturation has been observed [30]. In addition Shetty et al [37] have reported a negative role of androgens in the recovery of spermatogenesis in irradiated rats and in mice the sperm capacitation, acrosome reaction and fertilizing capacity are improved by estradiol and even more by phytoestrogens [38].

Today several experimental models of mice have been developed and shown helpful to clarify the estrogens role in vertebrates [39]. There is evidence from estrogen receptor gene knock out (ER α KO) mouse that estrogens are necessary for the achievement of fertility [40]. The male mice deficient in aromatase (ArKO) develop normally, the males are able to breed and to produce litters; nevertheless starting at the age of 5 months ArKO males start to have failures of spermatogenesis and by the age of one year all male mice are infertile [41].

In men it has been shown that the aromatase deficiency consecutive to a P450 aromatase gene mutation leads to sterility [42,43]; to our knowledge from the 6 reported (Table 1) cases 3 of them showed a low sperm counts number [44-46].

An inactivating mutation in the ER α gene (exon 2) has been reported by Smith et al [47] in an infertile man with a number of spermatozoa in the normal range although the viability was diminished. In addition a correlation between the amount of estradiol in the seminal plasma and the germ cell number has been shown [48] as well as a positive role for estradiol in stimulating the motility of human spermatozoa [49].

Durkee et al [50] have reported the existence of ER in human sperm and in addition, it has been demonstrated that the sperm membrane contains an estrogen receptor-related protein able to bind steroids (see for review [51]).

Obviously these membrane receptors would be connected with a signal transduction pathway involving quick answers such as the calcium channel and a calcium/calmoduline complex, known to be concerned for instance in sperm mobility and capacitation [52]. The demonstration of the production of nitric oxide in spermatozoa is now

considered as an alternative pathway for improving the sperm motility and capacitation [53]. Together with the existence of aromatase [21] the intracrine role of estrogens should be considered in spermatozoa.

Conclusion

Today it is clear that not only testicular somatic cells but also germ cells represent an additional source of estrogens in several species of mammals including man. Germ cells (both meiotic and post-meiotic cells) do not only produce estrogens but since they contain estrogen receptors that would explain part of the role (intracrine and / or paracrine) of estrogens in male germ cell development. The mechanisms of action of estrogens in the reproductive organs of the male (Figure 1) remain to be clarified as well as the regulation of the aromatase gene expression, not yet fully understood especially according to the testicular development. Furthermore one should kept in mind that not only rodent spermatozoa but ejaculated human spermatozoa express a functional aromatase and together with ER these data open new considerations about the role of estrogens all along the male genital tract and may be also in sperm fertilizing ability (Figure 2).

Acknowledgments

The authors express their warm thanks for Dr Jérôme Levallet who has been a pioneer in that field in my team.

References

- Carreau S, Genissel C, Bilinska B and Levallet J **The oestrogen sources in the testis and the reproductive tract of the male** *Inter J Androl* 1999, **22**:211-213
- Hess RA **Oestrogen in fluid transport in efferent ducts of the male reproductive tract** *Rev Reprod* 2000, **5**:84-92
- Carreau S **Germ cells: a new source of estrogens in the male gonad** *Mol Cell Endocrinol* 2001, **178**:65-72
- Carreau S, Bourguiba S, Lambard S, Galeraud-Denis I, Genissel C and Levallet J **Reproductive system: aromatase and estrogens** *Mol Cell Endocrinol* 2002, **193**:137-143
- O'Donnell L, Robertson KM, Jones ME and Simpson ER **Estrogen and spermatogenesis** *Endocr Rev* 2001, **22**:289-318
- Scobie GA, Macpherson S, Millar MR, Groome NP, Romana P and Saunders PTK **Human estrogen receptors: differential expression of ERalpha and beta and the identification of ERbeta variants** *Steroids* 2002, **67**:985-992
- Simpson ER, Mahendroo MS, Means GD, Kilgore MW, Hinshelwood MM, Graham-Lorence S, Amarneh B, It Y, Fisher CR, Dodson MM, Mendelson CR and Bulun SE **Aromatase cytochrome P450, the enzyme responsible for estrogen biosynthesis** *Endocr Rev* 1994, **15**:342-355
- Sebastian S and Bulun SE **A highly complex organization of the regulatory region of the human CYP19 (aromatase) gene as revealed by the human genome project** *J Clin Endocrinol Metab* 2001, **86**:460-4602
- Levallet J, Bilinska B, Mittre H, Genissel C, Fresnel J and Carreau S **Expression and immunolocalization of functional cytochrome P450 aromatase in mature rat testicular cells** *Biol Reprod* 1998, **58**:919-926
- Janulis L, Bahr JM, Hess RA, Jansen S, Osawa Y and Bunick D **Rat testicular germ cells and epididymal sperm contain active P450 aromatase** *J Androl* 1998, **19**:65-71
- Wiszniewska B **Primary culture of rat epididymal epithelial cells as a source of estrogen** *Andrologia* 2002, **34**:1-8
- Nitta H, Bunick D, Hess RA, Janulis L, Newton SC, Millette CF, Osawa Y, Shizuta Y, Toda K and Bahr JM **Germ cells of the mouse testis express P450 aromatase** *Endocrinology* 1993, **132**:1396-1401

13. Bilinska B, Schmalz-Fraczek B, Sadowska J and Carreau S **Immunolocalization of cytochrome P450 aromatase and oestrogen receptors α and β in bank vole testicular cells** *Acta Histochem* 2000, **102**:167-181
14. Kotula-Balak , Slomczynska M, Fraczek B, Bourguiba S, Tabarowski Z, Carreau S and Bilinska B **Complementary approaches demonstrate that cellular aromatization in the bank vole testis is related to photoperiod** *Eur J Histochem* 2003, **47**:55-62
15. Sipahutar H, Sourdain P, Moslemi S, Plainfossé B and Seralini GE **Immunolocalization of aromatase in stallion leydig cells and seminiferous tubules** *J Histochem Cytochem* 2003, **51**:311-318
16. Pereyra-Martinez AC, Roselli CE, Stadelman HL and Resko JA **Cytochrome P450 aromatase in testis and epididymis of male rhesus monkeys** *Endocrine* 2001, **16**:15-19
17. Carreau S **Paracrine control of human Leydig cell and Sertoli cell functions** *Folia Histochem Cytobiol* 1996, **34**:111-119
18. Chew PCT, Loganath A, Peh KL, Chow WP, Gunasegaram R and Ratman SS **Concentrations of intracellular sex steroids in human spermatozoa** *Arch Androl* 1993, **30**:165-170
19. Turner KJ, Macpherson S, Millar MR, McNeilly AS, Williams K, Cranfield M, Groome NP, Sharpe RM, Fraser HM and Saunders PTK **Development and validation of a new monoclonal antibody to mammalian aromatase** *J Endocrinol* 2002, **172**:21-30
20. Rago V, Bilinska B, Palma A, Ando S and Carpino A **Evidence of aromatase localization in cytoplasmic droplet of human immature ejaculated spermatozoa** *Folia Histochem & Cytobiol* 2003, **41**:23-28
21. Lambard S, Galeraud-Denis I, Bouraïma H, Bourguiba S, Chocat A and Carreau S **Expression of aromatase in human ejaculated spermatozoa: a putative marker of motility** *Molec Human Reprod* 2003,
22. Lanzino M, Catalano S, Genissel C, Ando S, Carreau S, Hamra K and McPhaul MJ **Aromatase messenger RNA is derived from the proximal promoter of the aromatase gene in Leydig, Sertoli, and germ cells of the rat testis** *Biol Reprod* 2001, **64**:1439-1443
23. Bourguiba S, Genissel C, Lambard S, Bouraïma H and Carreau S **Regulation of aromatase gene expression in leydig cells and germ cells** *J Steroid Biochem Molec Biol* 2003,
24. Shozu M, Zhao Y and Simpson ER **TGF β 1 stimulates expression of the aromatase (Cyp19) gene in human osteoblast and THP-1 cells** *Mol Cell Endocrinol* 2000, **160**:123-133
25. Genissel C, Levallet J and Carreau S **Regulation of cytochrome P450 aromatase gene expression in adult rat Leydig cells. Comparison with estradiol production** *J Endocrinol* 2001, **168**:95-105
26. Golovine K, Schwerin M and Vanselow J **Three different promoters control expression of the aromatase cytochrome p450 gene (cyp 19) in mouse gonads and brain** *Biol Reprod* 2003, **68**:978-984
27. Mowa CN and Iwanaga T **Expression of estrogen receptor-alpha and -beta mRNAs in the male reproductive system of the rat as revealed by in situ hybridization** *J Molec Endocrinol* 2001, **26**:165-174
28. Saunders PTK, Sharpe RM, Williams K, Macpherson S, Urquart H, Irvine DS and Millar M **Differential expression of oestrogen receptor α and β proteins in the testes and male reproductive system of human and non-human primates** *Mol Hum Reprod* 2001, **7**:227-236
29. Li H, Papadopoulos V, Vidic B, Dym M and Culty M **Regulation of rat testis gonocyte proliferation by platelet-derived growth factor and estradiol: identification of signaling mechanisms involved** *Endocrinology* 1997, **138**:1289-1298
30. Shetty G, Krishnamurthy H, Krishnamurthy HN, Bhatnagar S and Moudgal RN **Effect of estrogen deprivation on the male reproductive physiology of male and female primates** *J Steroid Biochem Molec Biol* 1997, **61**(3-6):26-30
31. Bilinska B, Schmalz-Fraczek B, Kotula M and Carreau S **Photoperiod-dependent capability of androgen aromatization and the role of estrogens in bank vole testis visualized by means of immunohistochemistry** *Mol Cell Endocrinol* 2001, **178**:189-198
32. Atanassova N, McKinnell C, Turner KJ, Walker M, Fisher JS, Morley M, Millar MR, Groome NP and Sharpe RM **Comparative effects of neonatal exposure of male rats to potent and weak (environmental) estrogens on spermatogenesis at puberty and the relationship to adult testis size and fertility: evidence for stimulatory effects of low estrogen levels** *Endocrinology* 2000, **141**:3898-3907
33. Ebling FJP, Brooks AN, Cronin AS, Ford H and Kerr JB **Estrogenic induction of spermatogenesis in the hypogonadal mouse** *Endocrinology* 2000, **141**:2861-2869
34. Pak TR, Lynch R and Tsai PS **Estrogen accelerates gonadal recrudescence in photo-regressed male Siberian hamsters** *Endocrinology* 2002, **143**:4131-4134
35. Bilinska B, Gancarczyk , Kotula-Balak M and Slomczynska M **Effect of 17 β -estradiol on testicular structure and immunoexpression of aromatase in immature bank vole testes** *Vlth International Aromatase Conference, Kyoto Abs PI-8* 2002, Oct 26-30
36. Pentakainen V, Erkkila K, Suomalainen L, Parvinen M and Dunkel L **Estradiol acts as a germ cell survival factor in the human testis** *J Clin Endocrinol Metab* 2000, **85**:2057-2067
37. Shetty G, Wilson G, Hardy MP, Niu E, Huhtaniemi I and Meistrich ML **Inhibition of recovery of spermatogenesis in irradiated rats by different androgens** *Endocrinology* 2002, **143**:3385-3396
38. Adeoya-Osiguwa SA, Markoulaki S, Pocock V, Milligan SR and Fraser LR **17 beta-estradiol and environmental estrogens significantly affect mammalian sperm function** *Human Reprod* 2003, **18**:100-107
39. Cooke HJ and Saunders PTK **Mouse models of male infertility** *Nature Rev Genetics* 2002, **3**:790-801
40. Couse JF and Korach KS **Estrogen receptor null mice : what have we learned and where will they lead us?** *Endocr Rev* 1999, **20**:358-417
41. Robertson KM, O'Donnell L, Jones MEE, Meachem SJ, Boon WC, Fisher CR, Graves KH, McLachlan RI and Simpson ER **Impairment of spermatogenesis in mice lacking a functional aromatase (Cyp 19) gene** *Proc Natl Acad Sc USA* 1999, **96**:7986-7997
42. Carani C, Qin K, Simoni M, Faustini-Fustini M, Serpente S, Boyd J, Korach KS and Simpson ER **Effect of testosterone and estradiol in a man with aromatase deficiency** *New Engl J Med* 1997, **337**:91-95
43. Morishima A, Grumbach MM, Simpson ER, Fisher C and Qin K **Aromatase deficiency in male and female siblings caused by a novel mutation and the physiological role of estrogens** *J Clin Endocrinol Metab* 1995, **80**:3689-3698
44. Carani C, Fabbri M, Zirilli L and Sgarbi I **Estrogen resistance and aromatase deficiency: clinical aspects in men** *J Soc Bi* 2002, **196**:245-248
45. Herrmann BL, Saller B, Janssen OE, Gocke P, Bockisch A, Sperling H, Mann K and Broecker M **Impact of estrogen replacement therapy in a male with congenital aromatase deficiency caused by a novel mutation of the CYP 19 gene** *J Clin Endocrinol Metab* 2002, **87**:5476-5484
46. Kottler ML, Piura M, Mittre H and Carreau S **Clinical findings in an adult man with a novel mutation in the aromatase gene** *Vlth International Aromatase Conference Abs PII-2-30* 2002 Kyoto 26-30 Oct
47. Smith EP, Boyd J, Frank GR, Takahashi H, Cohen RM, Specker B, Williams TC, Lubahn DB and Korach KS **Estrogen resistance caused by a mutation in the estrogen-receptor gene in a man** *New Engl J Me* 1994, **331**:1056-1061
48. Purvis K, Lindgren M, Cekan Z and Diczfalusy E **Indices of gonadal function in the human male. Seminal plasma levels of steroids in normal and pathological conditions** *Clin Endocrinol* 1975, **4**:247-248
49. Cheng CY and Boettcher B **The effects of steroids on the in vitro migration of washed human spermatozoa in modified tyrode's solution or in fasting human blood serum** *Fertil Steril* 1979, **32**:566-570
50. Durkee TJ, Mueller M and Zinaman M **Identification of estrogen receptor protein and messenger ribonucleic acid in human spermatozoa** *Amer J Obst Gyn* 1998, **178**:1288-1295
51. Luconi M, Forti G and Baldi E **Genomic and nongenomic effects of estrogens : molecular mechanisms of action and clinical implications for male reproduction** *J Steroid Biochem Molec Biol* 2002, **80**:369-381
52. Revelli A, Massobrio M and Tesarik J **Nongenomic actions of steroid hormones in reproductive tissues** *Endocr Rev* 1998, **19**:3-17
53. Herrero MB and Gagnon C **Nitric oxide : a novel mediator of sperm function** *J Androl* 2001, **22**:349-356