

RESEARCH

Open Access

# Functional expression and intracellular signaling of UTP-sensitive P2Y receptors in theca-interstitial cells

Francisco G Vázquez-Cuevas, Erika P Zárate-Díaz, Edith Garay and Rogelio O Arellano\*

## Abstract

**Background:** Purinergic receptors are expressed in the ovary of different species; their physiological roles remain to be elucidated. UTP-sensitive P2Y receptor activity may regulate cell proliferation. The aim of the present work was to study the functional expression of these receptors in theca/interstitial cells (TIC).

**Methods:** TIC were isolated by centrifugation in a Percoll gradient. P2Y receptors and cellular markers in TIC were detected by RT-PCR and Western blot. Intracellular calcium mobilization induced by purinergic drugs was evaluated by fluorescence microscopy, phosphorylation of MAPK p44/p42 and of cAMP response element binding protein (CREB) was determined by Western blot and proliferation was quantified by [<sup>3</sup>H]-thymidine incorporation into DNA.

**Results:** RT-PCR showed expression of p2y2r and p2y6r transcripts, expression of the corresponding proteins was confirmed. UTP and UDP, agonists for P2Y2 and P2Y6 receptors, induced an intracellular calcium increase with a maximum of more than 400% and 200% of basal level, respectively. The response elicited by UTP had an EC<sub>50</sub> of 3.5 +/- 1.01 μM, while that for UDP was 3.24 +/- 0.82 μM. To explore components of the pathway activated by these receptors, we evaluated the phosphorylation induced by UTP or UDP of MAPK p44 and p42. It was found that UTP increased MAPK phosphorylation by up to 550% with an EC<sub>50</sub> of 3.34 +/- 0.92 and 1.41 +/- 0.67 μM, for p44 and p42, respectively; these increases were blocked by suramin. UDP also induced p44/p42 phosphorylation, but at high concentrations. Phosphorylation of p44/p42 was dependent on PKC and intracellular calcium. To explore possible roles of this pathway in cell physiology, cell proliferation and hCG-induced CREB-phosphorylation assays were performed; results showed that agonists increased cell proliferation and prevented CREB-phosphorylation.

**Conclusion:** Here, it is shown that UTP-sensitive P2Y receptors are expressed in cultured TIC and that these receptors had the ability to activate mitogenic signaling pathways and to promote cell proliferation, as well as to prevent CREB-phosphorylation by hCG. Regulation of TIC proliferation and steroidogenesis is relevant in ovarian pathophysiology since theca hyperplasia is involved in polycystic ovarian syndrome. Purinergic receptors described might represent an important new set of molecular therapeutic targets.

## Background

Theca cells form a multilayer cover that surrounds the follicle beginning in its early developmental stages. The main physiological roles recognized for theca cells are the initial steps in the steroidogenic process; specifically, these cells convert acetate or cholesterol to androgens [1], which are secreted into the intra-follicular medium and

taken up by granulosa cells to serve as substrate for estrogen synthesis. In addition, theca cells could be an important signal integrator and regulator of aspects of follicular growth, because it represents the last follicular layer in contact with blood flow and receives chemical information from the peripheral nervous system [2].

Several studies in recent years indicate that the purinergic signaling system is functionally expressed in the ovary of several species [3] and represents another regulatory element in ovarian physiology; however, the physiological

\* Correspondence: arellano.ostoa@unam.mx

<sup>1</sup> Departamento de Neurobiología Celular y Molecular, Instituto de Neurobiología, Universidad Nacional Autónoma de México, Boulevard Juriquilla 3001, Juriquilla Querétaro, CP 76230, México  
Full list of author information is available at the end of the article

role of ATP in this context and its membrane receptors is unknown.

ATP is an important neurotransmitter in the peripheral nervous system [3], and nerve terminals from this system are potential sources for ATP release in the ovary. For example, the ovary is innervated by sympathetic terminals through the superior ovarian nerve and ovarian plexus [2,4]. It has been shown in other tissues that ATP is co-released with noradrenaline by sympathetic terminals and that it participates in several physiological events such as the induction and regulation of smooth muscle contraction [5] and the modulation of cardiac muscle excitation [6]. In addition, several cell types are able to release ATP in a basal manner and/or in response to different stimuli, such as mechanical stimulation, changes in pH, or hypotonic stress [7-9].

As a cellular messenger, ATP exerts its action through membrane receptors named P2, which are grouped into two subfamilies: P2X receptors that are cationic channels, and P2Y receptors that belong to the G protein-coupled receptor (GPCR) super family. In mammals, 8 subtypes of P2Y receptors have been described: 1, 2, 4, 6, and 11-14. Subtypes P2Y1, 2, 4, 6, and 11 are mainly coupled to  $G_{\alpha_q/11}$  proteins, and they activate phospholipase C (PLC) and consequently diacylglycerol and phosphoinositide- $Ca^{2+}$  turnover; subtypes 12-14, on the other hand, are coupled to  $G_{\alpha_{i0}}$  proteins that signal primarily by inhibiting adenylyl cyclase [10].

P2Y2, P2Y4, and P2Y6 form a subgroup of receptors sensitive to uridine nucleotides [11]; P2Y2 and P2Y4 show selectivity for nucleoside triphosphates, while P2Y6 prefers mainly nucleoside diphosphates, specifically UDP [12]. Uridine P2Y-activated receptors are involved in a broad variety of physiological processes such as cell proliferation, smooth muscle contraction, transmitter release, and others [3,10]. In the ovary, expression of UTP-sensitive P2Y receptors has been described in granulosa luteal cells [13,14], in the cumulus cell-oocyte complex [15], and in *Xenopus* ovarian follicles [16,17].

Recently, it was demonstrated that functional P2X7 receptors are expressed in mammalian TIC and can induce apoptotic cell death [18]. In the same study, it was also observed that the application of UTP evoked intracellular  $[Ca^{2+}]_i$  changes, suggesting that multiple P2 receptor subtypes are expressed in theca cells. Here, we studied this response in order to elucidate in more detail the molecular elements involved and the physiological implications of their activation. We found that uridine-sensitive P2Y2 and P2Y6 receptors are expressed in the TIC membrane and that P2Y activation promoted three important responses in these cells: 1) elicited  $Ca^{2+}$  mobilization from intracellular reservoirs, increasing the concentration of this important second messenger in the

cytoplasm; 2) increased cell proliferation through a mechanism dependent on the activation of protein kinase C (PKC) as well as MAPK p44 and p42, and; 3) down regulated hCG-dependent phosphorylation of CREB, an important element in steroidogenesis cascade control.

## Methods

### Materials

ATP, UTP, UDP, suramin, human chorionic gonadotropic hormone (hCG), porcine follicle stimulating hormone (FSH), and PPADS were purchased from Sigma Chemical Co. (St. Louis, MO), and staurosporin, wortmanin, and phorbol 12-myristate 13-acetate (PMA) were from Calbiochem (Gibbstown, NJ). DMEM-F12 medium, fetal bovine serum, antibiotic-antimycotic mix, and other cell culture reagents were from Gibco Invitrogen Co. (Grand Island, NY). Antibodies against mouse total or phosphorylated MAPK p44 and p42 and total or phosphorylated CREB were from Cell Signaling (Danvers, MA), and antibody against poly(ADP-ribose) polymerase-1 (PARP) was from Santa Cruz Biotechnology (Santa Cruz, CA). Oligonucleotides, reverse transcriptase, oligo dT, *taq* polymerase, and other molecular biology reagents were purchased from Invitrogen Co. (Carlsbad, CA), and Fluo4-AM was from Molecular Probes Invitrogen Co. (Eugene, OR). Automatic sequencing was done in the Molecular Biology Unit of the Instituto de Neurobiología, UNAM.

### Theca cell isolation and culture

Mouse theca/interstitial cells were purified by a discontinuous Percoll gradient [19,20]. Immature mice were used to avoid cultures enriched in luteal cells. Thus, intact 4- to 5-week-old mice of the strain C57 were sacrificed by cervical dislocation, a procedure approved by the ethical committee of Instituto de Neurobiología-UNAM. The ovaries were dissected and incubated in collagenase (100 mg/ml) for 20 min, and the ovarian epithelium was removed by passing the complete ovary repeatedly in and out through the orifice of a Pasteur pipette. Most granulosa cells were then eliminated by puncturing the isolated, epithelium-free ovaries with a fine hypodermic needle. The ovary, free of epithelium and most granulosa, was cut into fine pieces that were then incubated in a mix of collagenase (4 mg/ml), DNase I (10  $\mu$ g/ml), and BSA (10 mg/ml) for 30 min. The homogenate was fractionated on a discontinuous gradient: bottom layer 44% Percoll; top layer Percoll of density 1.055 g/ml. The cells were centrifuged for 20 min at  $400 \times g$ , and TIC were collected from the interface by aspiration, then cultured in DMEM-F12 medium containing 10% fetal bovine serum supplemented with antibiotic-antimycotic at 37°C in a humidified atmosphere with 5%  $CO_2$ . Cultures were maintained 48 h before using them in experiments, to allow proper

recovery after the isolation procedure. With this method we usually obtained  $1 \times 10^6$  cells per mouse; purity was assayed by RT-PCR and immunocytochemistry against CYP11A protein (> 96% of cells were positive). To exclude contamination with granulosa, the expression of the FSH receptor transcript and the responsiveness of CREB phosphorylation to hCG or FSH were assayed.

### Reverse Transcription Polymerase Chain Reaction

Total RNA of TIC cultures or from the indicated organ was purified using the guanidine isothiocyanate method [21]. First strand cDNA was synthesized using 2  $\mu$ g of DNase-treated RNA as template, 1 mg of oligo(dT), random hexamers, and reverse transcriptase.

The cDNA was used as template in a polymerase chain reaction to amplify cDNA fragments for  $\beta$ -actin, *p2y2r*, *p2y4r*, and *p2y6r* transcripts, and for *cyp11A*, *cyp17A*, *star*, and *fshr* as positive and negative theca cell markers, respectively. All the PCR programs started at 96°C for 3 min and finished at 72°C for 1 min. The amplification cycles consisted in 40 s at 96°C, 40 s at the specific annealing temperature for each primer set, and 40 s at 72°C.

The sequences of the oligonucleotides, the annealing temperatures, and the number of PCR cycles used were as follows: *p2y2r*, forward GGACGAACTGGGATA-CAAGTGT, reverse GTGGACTCTGTCCGTCTT-GAGT, annealing temperature 55°C, 30 cycles; *p2y4r*, forward GGGACTAACTGCAGGCAGAG, reverse GATACACATCAGGCCCGTCT, annealing temperature 60°C, 40 cycles; *p2y6r*, forward TTTCAAGCGACTGCT-GCTAA, reverse TGGCATAGAAGAGGAAGCGT, annealing temperature 55°C, 30 cycles; *cyp11A*, forward GCTGGAAGGTGTAGCTCAGG, reverse CACTGGT-GTGAACATCTGG, annealing temperature 55°C, 30 cycles; *cyp17A*, forward TGGTCGGCCCCAGATG-GTGA, reverse ATCTCGGGACTCCCCGTCTGT, annealing temperature 56°C, 30 cycles; *star*, forward AACAGGAAGGCTGGAAGAAG, reverse AGCAGC-CAAG TGAGTTTATGTC, annealing temperature 55°C, 30 cycles; *fshr*, forward TGGATGTCATCACTGGCTGT, reverse CAAATCTCAGTTCAATGGCG, annealing temperature 58°C, 30 cycles; and  $\beta$ -actin, forward GGGTCA-GAAGGATTCCCTATG, reverse GGTCTCAAACATGATCTGGG, annealing temperature 55°C, 25 cycles.

The amplified products were gel-isolated, phenol-chloroform purified, and subcloned into the pCR4-TOPO vector (Molecular Probes Invitrogen Co.). Their nucleotide sequences were confirmed by automatic sequencing.

### Fluorescence microscopy

Mouse ovarian TIC were grown on 12-mm diameter cover slides. Semi-confluent cultures were loaded for 15

min with 5 mM fluo-4/AM and 0.1% pluronic acid in Krebs solution (in mM: 150 NaCl, 1 KCl, 1 MgCl<sub>2</sub>, 1.5 CaCl<sub>2</sub>, 4 glucose, 10 HEPES, and 0.05% BSA, pH 7.4). The cells were washed with Krebs solution for 10 min to eliminate excess dye and then placed in a constant-flow recording chamber that allowed them to be visualized with an inverted fluorescence microscope (Olympus IX70, Melville, NY). Drugs were applied by superfusion and responses were recorded with an Evolution QEi camera (Media Cybernetics, Bethesda, MD). Sequences of images were analyzed using the Image-Pro Plus software (Media cybernetics, Bethesda, MD) and Imagenes software, a program developed specifically for this analysis (Dr. Ivan Terol, CIDETEQ, México). In the Ca<sup>2+</sup>-free Krebs solution, CaCl<sub>2</sub> was replaced by 3 mM MgCl<sub>2</sub>.

### Western blot

For MAPK p42 and p44 or CREB phosphorylation experiments, cultured TIC ( $5 \times 10^5$ ) were harvested 24 h before the experiment to reduce serum-dependent kinase activity. After that, cells were stimulated with the indicated drugs, scraped in Laemmli buffer, and boiled for 5 min. For electrophoresis, samples were fractionated in a 10% SDS-polyacrylamide gel and transferred to a nitrocellulose membrane (BioRad, Hercules, CA). Membranes were blocked for 1 h at room temperature in 150 mM NaCl, 20 mM Tris, pH 7.4, and 0.1% Tween 20 (TBS-T) containing 5% nonfat dry milk and then incubated overnight at 4°C with the appropriate rabbit primary antibody (1:1000) directed against the phosphorylated form of MAPK p44 and p42 or CREB (Cell signaling, Danvers, MA). After washing with TBS-T, membranes were incubated 1 h at 37°C with HRP-conjugated goat anti-rabbit antibody (Zymed, Invitrogen Co., Grand Island, NY) in TBS-T. The immunoreactive proteins were detected by chemiluminescence, and images were analyzed by pixel density with ImageJ Software (NIH, USA); the results were expressed in terms of optic density normalized against the basal condition, a parameter that is proportional to the change in protein phosphorylation. To analyze total p44/p42 or other load controls, such as PARP, the same membranes were incubated for 30 min in stripping solution (50 mM, Tris pH 6.8, 100 mM  $\beta$ -mercaptoethanol, and 2% SDS) at 55°C, washed twice with TBS-T, and then reprobed with a primary antibody against the indicated protein.

### Immunoprecipitation

TIC ( $1 \times 10^6$ ) were scraped in ice-cold TNTE buffer (Tris-HCl pH 7.4, 150 mM NaCl, 50 mM and 0.5 mM EDTA) containing 5% Triton X-100 and a protease inhibitor cocktail (Roche Co., Basel, Switzerland); the lysate was centrifuged for 10 min at 14,000 rpm at 4°C, and the soluble fraction was incubated overnight with 3  $\mu$ l of anti-

P2Y6 antibody (Alomone, Jerusalem, Israel). After that, 50  $\mu$ l of protein G agarose was added to the lysate and incubated for 1 h at room temperature; the agarose beads were washed 3 times with TNTE containing 1% Triton X-100 and protease inhibitors, resuspended in Laemmli buffer, boiled for 5 min, and analyzed by Western blot.

### Proliferation assay

Cell proliferation was analyzed using [ $^3$ H]-thymidine incorporation. For this, cells ( $10^4$ ) were cultured in 48-well plates and after 48 h of culture, they were harvested and incubated for 24 h in serum-free DMEM-F12 media; then the culture medium was changed to DMEM-F12 with 0.1% fetal bovine serum containing the experimental treatment (UTP, UDP, or ATP at the indicated concentration). Then cultures were incubated for another 48 h, with the addition of 1  $\mu$  Ci/well of [ $^3$ H]-thymidine after the first 24 h. At the end of the incubation, each well was washed 3 times with 5% trichloroacetic acid, and then the cells were lysed by addition of 250  $\mu$ l of boiling 250 mM NaOH, incubated 5 min, and transferred to vials containing 5 ml of scintillation liquid. Samples were counted in a scintillation counter.

### Statistical analysis


All data are expressed as mean  $\pm$  S.E.M. Statistical analysis was performed using GraphPad Prism (La Jolla, CA) software. The means of two groups were compared using a Student's t-test. ANOVA was used to compare several groups, and differences were considered to be significant at  $p < 0.05$ .

## Results

### Theca cell identity and expression of P2Y2, P2Y4, and P2Y6 receptors

TIC were isolated, and their identity was confirmed by RT-PCR amplification of *cyp11A*, *cyp17A*, and *star* transcripts as specific markers for theca cells, and of FSH receptor (*fshr*) transcripts as indicator of a possible contamination with granulosa cells; the  $\beta$ -actin transcript was used as a control housekeeping gene (Figure 1A). The results showed that TIC cultures were positive for *cyp11A*, *cyp17A*, and *star* expression, but they did not express the FSH receptor, demonstrating that the isolated cells were mainly of the thecal/interstitial type and were essentially free of granulosa cells.

This conclusion was strengthened with data obtained in functional experiments. For this, TIC cells were stimulated by 2 IU hCG or 1 ng/ml FSH, and CREB phosphorylation was evaluated. It is well established that gonadotropin receptors exert their actions by coupling to G proteins, increasing cAMP synthesis that, in consequence, promotes CREB-phosphorylation [22,23]. It was found that in TIC cultures, CREB protein-phosphoryla-



tion (43-kDa band) was increased 6-fold by hCG stimulation, whereas FSH did not induce any change (Figure 1B), thus supporting the idea that these cultures contained mainly theca/interstitial cells.

To study the expression of *p2y2r*, *p2y4r*, and *p2y6r* transcripts, RNA from TIC was reverse transcribed, and then PCR was carried out with specific oligonucleotides for each receptor subtype. RNA samples from ovary, brain, and heart were also analyzed as controls. As shown in Figure 1A, a *p2y2r* fragment of 1032 bp and a *p2y6r* fragment of 257 bp were amplified from the cDNA of all tissues tested. However, the *p2y4r* fragment of 575 bp was only amplified from the whole ovary and brain cDNA. In all the assays, control amplifications without RT or without cDNA template did not produce any PCR product (Figure 1A). The amplified fragments were cloned into the pCR4-TOPO vector, sequenced, and analyzed in BLAST, and the fragments were identical to the reported sequences from mouse (NM008773 for *p2y2r*, NM020621 for *p2y4r*, and NM183168 for *p2y6r*).

These RT-PCR results indicated that TIC might express P2Y2 and P2Y6 receptors. In order to detect the protein, Western blot was performed from homogenates to detect P2Y2; to detect P2Y6 receptor it was necessary to perform immunoprecipitation followed by Western blot,

which suggested a low expression level of this receptor (Figure 1C). P2Y2 was detected as a band of 58 kDa, a major band near 70 kDa, and a fainter band of 45 kDa. P2Y6 was detected as three bands with molecular weights of approximately 45, 40, and 37 kDa. In the latter case, the IgG heavy chain interfered with the immunoreactive bands corresponding to the receptor. However, all bands observed match the molecular weights reported previously for both receptor types [e.g., [24,25]].

#### UTP- and UDP-induced increase of intracellular $\text{Ca}^{2+}$ concentration in TIC


Functional responses of P2Y receptors were studied by applying ATP, UTP, or UDP to TIC and monitoring the changes in intracellular calcium concentration ( $[\text{Ca}^{2+}]_i$ ) using fluorescence microscopy of Fluo-4-AM loaded-cells. In all cases, 25 to 40 cells from 3 independent cultures (each from 2 donors) were analyzed (Figure 2).

Figure 2A shows a typical response elicited by 100  $\mu\text{M}$  ATP. At the highest concentration tested (1 mM), ATP elicited a  $[\text{Ca}^{2+}]_i$  increase of  $458 \pm 18\%$  compared with the basal level (Figure 2B, left panel); this increase was monotonic (Figure 2B, right panel), dose-dependent, and had an  $\text{EC}_{50}$  of  $6.5 \pm 1.0 \mu\text{M}$ . In cells from the same cultures, UTP also induced a dose-dependent response with an  $\text{EC}_{50}$  of  $3.5 \pm 1 \mu\text{M}$  and a maximal increase of  $437 \pm 12\%$  (Figure 2C, left panel). As illustrated in the right panel, the increase generated by UTP had a similar time-course to that elicited by ATP.

Three types of P2Y receptors sensitive to UTP have been described: P2Y2, P2Y4, and P2Y6 receptors. UDP is a more potent agonist for P2Y6 receptors than UTP or ATP; thus, in order to detect a possible participation of P2Y6, TIC were tested with UDP. This nucleotide elicited responses with an  $\text{EC}_{50}$   $3.2 \pm 0.8 \mu\text{M}$ ; however, the maximal response reached was only  $210 \pm 5.4\%$  (Figure 2D, left panel). Furthermore, the  $[\text{Ca}^{2+}]_i$  increase in response to UDP consistently showed an oscillating time-course (Figure 2D, right panel), different from that observed with ATP or UTP.

In the absence of extracellular  $\text{Ca}^{2+}$  ( $\text{Ca}^{2+}$ -free solution), responses to either UTP or UDP were not abolished (Figure 2E). Nevertheless, maximal responses generated by UTP averaged  $366 \pm 13\%$ , significantly less ( $P < 0.01$ ) than those observed in normal Krebs solution ( $437 \pm 12\%$ ). The  $\text{EC}_{50}$  obtained for UTP in  $\text{Ca}^{2+}$ -free solution was  $6.2 \pm 0.9 \mu\text{M}$  and was not significantly different from that obtained in normal Krebs. For UDP, similar findings were observed: the maximal response reached  $230 \pm 15\%$  and had an  $\text{EC}_{50}$  of  $4.9 \pm 0.6 \mu\text{M}$ ; neither parameter differed significantly from that in normal Krebs.

This suggested that extracellular  $\text{Ca}^{2+}$  was not the major source of the  $[\text{Ca}^{2+}]_i$  increase produced in TIC by



UTP or UDP; more probably, this increase came from intracellular reservoirs via  $\text{IP}_3$  synthesis, as shown in other cell systems.

#### UTP-induced activation of p44 and p42 MAPK

In order to study the signaling pathway involved in the UTP and UDP activation of P2Y receptors in TIC, phosphorylation of the p44 and p42 MAPK proteins was evaluated (in at least 3 experiments, each in duplicate, carried out on independent cultures derived from pooled ovaries of 2-3 mice). For these experiments, UTP was used as a specific agonist of the P2Y receptor subtypes studied. It


was observed that UTP induced MAPK phosphorylation in a dose-dependent manner with an  $EC_{50}$  of  $3.3 \pm 0.9$  and  $1.4 \pm 0.7 \mu\text{M}$  for p44 and p42, respectively; maximal increases of  $541 \pm 25.6\%$  and  $461 \pm 34.8\%$ , respectively, were observed by applying  $100 \mu\text{M}$  UTP (Figure 3A). The time course of this effect was studied by applying  $10 \mu\text{M}$  UTP and measuring p44 and p42 MAPK phosphorylation at different times. The results indicated that maximal phosphorylation occurred at 5 min of stimulation, and then it decreased slowly, returning to near-basal levels about 30 min after UTP addition (Figure 3B).

Because it has been shown consistently that UDP acts more potently on P2Y6 receptors [11,12], its ability to promote p44 and p42 MAPK phosphorylation was tested. In experiments similar to those presented above for UTP,  $100 \mu\text{M}$  UDP was less potent and induced only modest responses of  $199 \pm 43\%$  and  $158 \pm 15\%$  for p44 and p42, respectively, compared to the basal level; the effect increased to  $364 \pm 63\%$  and  $349 \pm 95\%$ , respectively, with


$1 \text{ mM}$  UDP (Figure 3C). The time course of p44/p42 phosphorylation induced by  $1 \text{ mM}$  UDP was similar to that elicited by  $100 \mu\text{M}$  UTP (Figure 3D). In addition, the p44 and p42 MAPK phosphorylation induced by  $10 \mu\text{M}$  UTP was antagonized by suramin with an  $IC_{50}$  of  $84.3 \pm 10.2 \mu\text{M}$  (Figure 4A); suramin is a potent antagonist of P2Y2 receptors but is only a weak antagonist of P2Y6 [11]. Conversely, PPADS up to  $600 \mu\text{M}$ , a drug that antagonizes mainly the P2Y6 receptor [11], had no effect on UTP-induced MAPK phosphorylation (Figure 4B). These results suggested that P2Y6 is not a major participant in the phosphorylation of MAPK; in consequence, the following experiments focused on defining the role of the P2Y2 receptor in the purinergic response.

#### UTP-induced p44 and p42 MAPK phosphorylation is dependent on PKC activation


UTP-dependent p44/p42 MAPK phosphorylation might be elicited via either of two main mechanisms: 1) transactivation of EGF receptors as has been demonstrated, for example, in salivary gland cells [26] or 2) by activation of kinases through its canonical pathway. However, EGF transactivation requires long-term (over 1 h) stimulation with the agonist, and activation of p44/p42 MAPK in TIC was generated even with short incubations; thus, it was decided to analyze first the role of down stream kinases. Two of the main candidate protein kinases, PKC and PI3K [27,28], were blocked using specific pharmacological tools. PI3K activation was blocked by preincubation for 30 min with the selective inhibitor  $100 \text{ nM}$  wortmanin (Figure 5A) or with  $1 \mu\text{M}$  LY294002 (not shown) before stimulating the cells with  $10 \mu\text{M}$  UTP; under these condi-



**Figure 3** D-R relationships and time course for the phosphorylation response elicited by UTP or UDP on p44 and p42 MAPK. (A) Phosphorylated isoforms of p44 (black dots) or p42 (white dots) were immunodetected (Western blot) in samples of theca cells that were stimulated with various UTP concentrations. The band intensities (OD) are expressed as a percent of the basal level, and the values were used to plot the D-R relationship. (B) Time course of the p44 and p42 phosphorylation process elicited by  $10 \mu\text{M}$  UTP. (C) D-R for phosphorylation elicited by UDP in experiments similar to those in (A). (D) Time course of p44 and p42 phosphorylation elicited by  $1 \text{ mM}$  UDP. Data points are the mean  $\pm$  S.E.M. of three independent experiments.




**Figure 4** p44 and p42 MAPK phosphorylation induced by UTP is blocked by suramin but not by PPADS. Cultures of mouse theca cells were incubated for 30 min with suramin (A) or PPADS (B) at the indicated concentrations; subsequently,  $10 \mu\text{M}$  UTP was applied for 5 min, and p44 (black dots) and p42 (white dots) MAPK phosphorylation was detected by Western blot. Data points are the mean  $\pm$  S.E.M. of three independent experiments.



tions, neither inhibitor had any effect on p44 or p42 MAPK phosphorylation ( $n = 3$  independent cultures).

To study the possible participation of PKC, TIC cultures were preincubated (30 min) with 250 nM staurosporine and then tested with 10  $\mu$ M UTP. Staurosporine treatment blocked completely the UTP-stimulated p44 and p42 phosphorylation (Figure 5A), strongly suggesting that phosphorylation was dependent on PKC. To support this idea, experiments were carried out in which PKC activity was downregulated by long-term incubations with phorbol-12-myristate-13-acetate (PMA) [29]. Thus, TIC were pretreated for 18 h with 1  $\mu$ M PMA, which did not affect the basal levels of phosphorylated p44 or p42 proteins; cells were then stimulated with 10  $\mu$ M UTP. Under these conditions, PMA preincubation reduced p44 and p42 MAPK phosphorylation induced by UTP from a maximal response without PMA of  $347 \pm 58\%$  and  $299 \pm 56\%$  for p44 and p42, respectively, to  $164 \pm 16\%$  and  $210 \pm 43\%$  ( $p < 0.05$ ) (Figure 5B). These results indicate that PKC is the main kinase responsible for the UTP-induced activation of the p44 and p42 proteins.

To test the role of intracellular  $Ca^{2+}$  during p44 and p42 MAPK phosphorylation, cell cultures were preincubated (90 min) with 10  $\mu$ M BAPTA-AM to load the cells intracellularly with this  $Ca^{2+}$  chelator. Fluorescence microscopy confirmed that this treatment prevented the calcium increase induced by UTP. In BAPTA-loaded TIC, phosphorylation of MAPK elicited by UTP was strongly inhibited (Figure 6); thus, in control conditions, UTP elicited a phosphorylation increase of  $384 \pm 53$  and  $289 \pm 55\%$  for p44 and p42, respectively, while UTP-stimulated BAPTA-




loaded cells showed significantly ( $p < 0.05$ ) lower phosphorylation increases of only  $171 \pm 40$  and  $116 \pm 16\%$ , respectively. This result indicates that phosphorylation was a  $Ca^{2+}$ -dependent process and provides evidence for PKC participation.

#### Evidence that suggests a role for purinergic signaling in TIC physiology

Cell proliferation is a consequence of purinergic stimulation in various cell systems [3]; here we asked whether or not P2Y stimulation of TIC induced their proliferation. For this, cell cultures were stimulated with different concentrations of UTP, ATP, or UDP; cell proliferation was estimated using [ $^3$ H]-thymidine incorporation and compared with that elicited by 10% FBS (Figure 7 shows four independent determinations, each in triplicate, from cultures where the ovaries of 2-3 mice were pooled).

The results indicated that ATP, UTP, or UDP increased proliferation. Incubation with 10% FBS increased [ $^3$ H]-thymidine incorporation to  $277 \pm 23\%$  of the basal level, and a similar increase was induced by 10  $\mu$ M ATP ( $232 \pm 33\%$ ) but not by 100  $\mu$ M ATP ( $135 \pm 13\%$ ). Also, 100  $\mu$ M UTP or UDP induced robust responses of  $205 \pm 18\%$  and  $221 \pm 17\%$ , respectively, while at 1 mM they generated



**Figure 7 Proliferative response and inhibition of hCG-induced CREB phosphorylation by P2Y agonists in mouse theca cells.** Theca cells were stimulated for 48 h with UTP, UDP, or ATP using the indicated concentrations (log M); controls corresponded to cells incubated in media supplemented with 0.1% serum (basal) or with 10% FBS (S). Cell proliferation was evaluated by  $^3\text{H}$ -thymidine incorporation, data are the mean  $\pm$  S.E.M. of 4 independent determinations. B) Time-course of CREB phosphorylation in TIC induced by 2 IU hCG; the 43-kDa band was quantified as in Figure 1B showing that maximal effect was reached in 10-15 min of incubation. C) CREB-phosphorylation elicited by 2 IU hCG in TIC incubated for 10 min in basal conditions and in the presence of 0, 10, or 100  $\mu\text{M}$  UTP; the 43-kDa band was quantified as in B), data are from 3 independent experiments in triplicate. (\* $p < 0.05$  vs. basal, one-way ANOVA and Bonferroni's test).

increases of  $216 \pm 37\%$  and  $183 \pm 23\%$ , respectively. The results strongly suggested that P2Y receptors were able to activate a proliferative response in TIC.

The possibility of regulatory cross talk between P2Y2 and LH receptors was also examined. First, it was shown that TIC responded to a 2 IU hCG stimulus by increasing CREB phosphorylation, and this response reached a maximum in 10 min (figure 7B). This observation demonstrated that the LH receptor efficiently activated the cAMP pathway in TIC cultures (see figure 1B). It is well established that the cAMP/PKA/CREB pathway participates in the canonical signal transduction cascade for regulation of androgen biosynthesis through the LH receptor [22].

With this rationale, cultures of TIC were incubated in 0, 10, or 100  $\mu\text{M}$  of UTP for 10 min, then 2 IU hCG were added for 10 min more, cell lysates were obtained, and CREB phosphorylation was evaluated by Western blot as an indicator of LH receptor activity. It was found that UTP, either 10 or 100  $\mu\text{M}$ , completely blocked CREB phosphorylation (figure 7C), strongly suggesting that P2Y2 receptor stimulation inhibited the cAMP pathway

activated by LH. This raises the possibility that the purinergic system participates in regulating the physiological actions promoted by LH, for example, the steroid hormones synthesis pathway.

## Discussion

It has been recognized that neurotransmitters might play distinct, regulatory roles in ovarian physiology; for example, it has been proposed that, in addition to the regulatory actions of gonadotropins, the activity of sympathetic fibers that innervate the ovary controls different aspects of ovarian function, such as steroidogenesis, folliculogenesis, and ovulation [e.g., [30,2,31,32]]; most of these studies have examined the role of the catecholaminergic system and specifically, of norepinephrine [30-32], but there is also a great deal of important evidence for participation of an ovarian cholinergic system [33,34]. Although knowledge about the purinergic system in the ovary is scarce, it is well established that ATP and norepinephrine are co-released at similar concentrations from sympathetic terminals in many cell systems [5,7], and release of ATP by the oocyte has already been documented in other species [8,16]. Thus, the study in the ovary of the molecular components expressed and cellular mechanisms activated by the purinergic system will be of importance to understand the possible role of ATP in ovarian physiology and pathology. Here, we present clear evidence of functional expression of UTP-sensitive P2Y receptors in TIC cultures, suggesting a role for these receptors in ovarian physiology.

RT-PCR and Western blot studies indicated that cultured TIC express P2Y2 and P2Y6 receptors. In functional experiments, UTP and UDP, specific agonists for P2Y2 and P2Y6, respectively, induced robust  $\text{Ca}^{2+}$  signals in normal Krebs or in  $\text{Ca}^{2+}$ -free solution, which indicated that the nucleotides promoted the response mainly through  $\text{Ca}^{2+}$  release from intracellular reservoirs, in agreement with the canonical  $\text{G}\alpha\text{q}$ -PLC pathway for these receptors [10]. UTP or high concentrations of UDP (1 mM) also induced the phosphorylation of MAPK p44 and p42; at high concentrations, UDP acted principally on the P2Y2 receptor, since P2Y6 is stimulated by UDP in the low  $\mu\text{M}$  range [11]. Phosphorylation of MAPK was inhibited by suramin, a potent antagonist for P2Y2 and weak for P2Y6, but it was not affected by PPADS, which is inactive toward P2Y2 but able to antagonize P2Y6 activation [11]. Taken together, our data indicated a main role of the P2Y2 receptor in MAPK activation. There is ample evidence that these protein kinases are involved in the proliferative phenomenon activated by G protein-coupled receptors in various cell systems [e.g., [27,28]]; in addition, p44 and p42 MAPK activation dependent on P2Y2 or P2Y6 receptors has been described, e.g., in granulosa-luteal cells [14], glioma cells [35], and embryonic



stem cells [36]. Staurosporin or long-term (18 h) incubation with PMA blocked UTP-induced p44 and p42 MAPK phosphorylation. In addition, p44 and p42 MAPK phosphorylation was blocked in BAPTA-loaded cells, strongly suggesting that a calcium-dependent PKC participates in this response.

Activation of MAPK p44 and p42 is directly related to induction of cell proliferation [37]. Our results demonstrated that UTP and UDP induced a robust proliferative response similar to that of 10% FBS used as positive control. ATP induced a proliferative response at 10  $\mu$ M, but no effect was observed with higher concentrations. This supports the idea that P2Y2 is the main receptor involved in the response, but an ancillary participation of P2Y6 cannot yet be excluded. The regulation of theca cell proliferation is relevant during folliculogenesis [1], and it might be involved in pathological processes, such as the altered androgen-estrogen balance associated with polycystic ovary syndrome, a common disease characterized by uncontrolled theca cell proliferation [38]. In this context, purinergic signaling can activate a feedback mechanism by inducing a proliferative or an apoptotic response in TIC.

ATP actions to stimulate TIC proliferation through P2Y2 (and P2Y6) receptor activation should be taken into account, together with the effects described for other neurotransmitters that seem to regulate specific processes in the ovary. For example, previous evidence showed that human granulosa-luteal cells express M1 and M5 muscarinic receptors [34] as well as P2Y2 purinergic receptors [13,14]; stimulation of either system by acetylcholine or ATP can promote granulosa-luteal cell proliferation. Stimulation of  $\beta$ -adrenergic receptors also modulates steroidogenic activity and ovulation [31] and, given that neurotransmitters released from catecholaminergic terminals might include ATP, it would be of interest to know the effect of activating purinergic receptors in these processes.

The results also showed that P2Y2 receptor activation had an important effect on the LH signaling pathway. It has been shown before that LH induces CREB phosphorylation [22,39] and that expression of a dominant negative CREB variant was enough to block androgen biosynthesis in rat TIC cells [22]. We observed that preincubation with UTP (10 or 100  $\mu$ M), completely blocked the hCG-induced CREB phosphorylation, which suggests that the purinergic system potently modulates LH-activated pathways, an action that might have important consequences in ovarian theca physiology.

Is well known that during folliculogenesis LH exerts regulatory actions beginning around the formation of early secondary follicles, which is concurrent with theca layer differentiation; from this stage throughout folliculogenesis up to ovulation, LH is the main regulator of theca

layer development, because it controls the steroidogenesis process [40]. However, during this period, important phenomena such as follicular selection or dominance processes cannot be explained solely by LH action; paracrine and autocrine follicular molecules seem to be essential for the final outcome [41]. It is possible that P2Y2 activation represents one of the mechanisms by which LH regulates the cohort of follicles that will or will not become dominant. Thus, the process of purinergic regulation demonstrated here might be involved in maintaining the proper balance between the rate of cell division and death in the ovary, and in essential physiological actions such as steroidogenesis, functioning as a local, fine-tuning modulator to complement the systemic control exerted by hormones and nervous system afferents. Hence, purinergic regulation is a potential therapeutic target in ovarian pathologies where proliferation or the steroidogenesis processes are affected.

Specifically in regulating the balance between theca proliferation and death, our data suggest that activation of the purinergic system by ATP could have dual effects on theca cell physiology; i.e., depending on the concentration, ATP might induce: 1) apoptotic cell death through P2X7 receptors (18) and 2) cell proliferation through P2Y2/P2Y6 receptors, as shown here. This is similar to what has been demonstrated in other systems in which the cells seem to co-express multiple purinergic receptor subtypes, leading to activation of multiple signaling pathways. For example, macrophages express a variety of P2X and P2Y purinergic receptors, and their activation modulates diverse physiological process such as apoptosis, activation of cell proliferation pathways, or activation of the inflammatory response machinery [42,43]. The final physiological outcome of the effect exerted by ATP in a given process will be determined by several factors including, for example, the purinergic receptor affinities, source and availability of ATP, ecto-ATPase activity, and also cross-talk between different G protein-coupled receptor types or subunits of receptor channels [see e.g., [17]]. In this context, it is important to mention that high concentrations of ATP (0.1 - 1 mM), but not of UTP, were consistently unable to increase cell proliferation, which might be a result of P2X7 receptor activation that can induce apoptotic cell death [18], among other possibilities, such as a regulatory effect of ATP on P2Y2/P2Y6 receptor function. Distinguishing among the various possibilities will require further analysis of the functional interaction among the different P2 receptors expressed in the ovarian theca.

Data presented in the present work are the first evidence that UTP-sensitive P2Y receptors are expressed and functional in theca cells. Although extensive studies are necessarily to establish with detail the main physiological activities, experimental data suggested these

receptors have a role in p44/p42 MAPK phosphorylation, proliferation increase, and cross talk with LH-activated pathways. These observations raise the possibility that the purinergic signaling system represents an important physiological regulator of theca cells.

## Conclusion

In summary, it was shown here that TIC express functional P2Y2 and P2Y6 receptors, which, when stimulated, induce a Ca<sup>2+</sup>-dependent proliferative response mediated through PKC activation and phosphorylation of the p42 and p44 MAPK proteins. P2Y receptor stimulation also regulates hCG-dependent CREB phosphorylation, suggesting interactions between functional pathways. Molecular components of purinergic transmission systems represent new molecular targets that must be characterized in the context of ovarian pathophysiology.

## List of abbreviations

P2Y: Purinergic G protein-coupled receptors; MAPK: Mitogen-activated protein kinases; PPADS: pyridoxal phosphate-6-azo (benzene-2,4-disulfonic acid) tetrasodium salt; BAPTA: 1,2-bis(o-aminophenoxy)ethane-N, N, N', N'-tetraacetic acid; CREB: cAMP response element binding protein; TIC: Theca interstitial cells; PMA: Phorbol 12-myristate 13-acetate.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

FGVC and ROA conceived and designed the study, carried out experiments, performed the data analysis, and drafted the manuscript; EPZD and GE performed experiments and participated in the analysis of data. All authors read and approved the final manuscript.

## Acknowledgements

This work was supported by: CONACYT-México No. 82340 to R.O.A.; PAPIIT-UNAM, México, No. IN208209 to F.V.-C. and IN214409 to R.O.A. We are grateful to Dr. Dorothy D. Pless for help with preparation of the manuscript. We also thank Mr. Horacio Ramírez Leyva, M. en C. Leonor Casanova, M.V.Z. Martín García Servín, Dr. Anaid Antaramian, Ing. Nydia Hernández Rios, Ing. Ramón Martínez Olvera, and M. en C. Alberto Lara Ruvalcaba for their expert technical assistance.

## Author Details

Departamento de Neurobiología Celular y Molecular, Instituto de Neurobiología, Universidad Nacional Autónoma de México. Boulevard Juriquilla 3001, Juriquilla Querétaro, CP 76230, México

Received: 22 March 2010 Accepted: 14 July 2010

Published: 14 July 2010

## References

1. Bjersing L: **Maturation, morphology and endocrine function of the follicular wall in mammals.** In *The mammalian ovary. Comparative Biology* Edited by: Jones RE. New York: Raven Press; 1978:181-214.
2. Burden HW: **The adrenergic innervation of mammalian ovaries.** In *Catecholamines as hormone regulators* Edited by: Ben-Jonathan N, Bahr JM, Weener R. New York: Raven Press; 1985:261-278.
3. Burnstock G: **Physiology and pathophysiology of purinergic Neurotransmission.** *Physiol Rev* 2007, **87**:659-797.
4. Owman C, Rosengren E, Sjöberg NO: **Adrenergic innervation of the human female reproductive organs: a histochemical and chemical investigation.** *Obstet Gynecol* 1967, **30**:763-773.
5. Westfall DP, Todorov LD, Mihaylova-Todorova ST: **ATP as a cotransmitter in sympathetic nerves and its inactivation by releasable enzymes.** *J Pharmacol Exp Ther* 2002, **303**(2):439-444.
6. Sesti C, Broekman MJ, Drosopoulos JH, Islam N, Marcus AJ, Levi R: **Ectonucleotidase in cardiac sympathetic nerve endings modulates ATP-mediated feedback of norepinephrine release.** *J Pharmacol Exp Ther* 2002, **300**:605-611.
7. Bodin P, Burnstock G: **Purinergic signalling: ATP release.** *Neurochem Res* 2001, **26**(8-9):959-969.
8. Saldaña C, Garay E, Rangel GE, Reyes LM, Arellano RO: **Native ion current coupled to purinergic activation via basal and mechanically induced ATP release in *Xenopus* follicles.** *J Cell Physiol* 2009, **202**:814-821.
9. Saldaña C, Vázquez-Cuevas FG, Garay E, Arellano RO: **Epithelium and/or theca are required for ATP-elicited k<sup>+</sup> current in follicle-enclosed *Xenopus* oocytes.** *J Cell Physiol* 2005, **218**:355-365.
10. Abbracchio M, Burnstock G, Boeynaems JM, Barnard EA, Boyer JL, Kennedy C, Knight G, Fumagalli M, Gachet C, Jacobson K, Weisman G: **International Union of Pharmacology LVIII: Update on the P2Y G protein-coupled nucleotide receptors: from molecular mechanisms and pathophysiology to therapy.** *Pharmacol Rev* 2006, **58**:281-341.
11. von Kugelgen I, Wetter A: **Molecular pharmacology of P2Y-receptors.** *Naunyn-Schmiedeberg's Arch Pharmacol* 2000, **362**:310-323.
12. Nicholas RA, Watt WC, Lazarowski ER, Li Q, Harden K: **Uridine nucleotide selectivity of three phospholipase C-activating P2 receptors: identification of a UDP-selective, a UTP-selective, and an ATP- and UTP-specific receptor.** *Mol Pharmacol* 1996, **50**:224-229.
13. Tai CJ, Kang SK, Choi KC, Tzeng CR, Leung PC: **Antigonadotropic action of adenosine triphosphate in human granulosa-luteal cells: Involvement of protein kinase Ca.** *J Clin Endocrinol Metab* 2001, **86**:3237-3242.
14. Tai CJ, Kang SK, Choi KC, Tzeng CR, Leung PC: **Adenosine triphosphate activates mitogen-activated protein kinase in human granulosa-luteal cells.** *Endocrinology* 2001, **142**:1554-1560.
15. Arellano RO, Martínez-Torres A, Garay E: **Ionic currents activated via purinergic receptors in the cumulus cell-enclosed mouse oocyte.** *Biol Reprod* 2002, **67**:837-846.
16. Arellano RO, Garay E, Miledi R: **Cl<sup>-</sup> currents activated via purinergic receptors in *Xenopus* follicles.** *Am J Physiol* 1998, **274**(2 Pt 1):C333-C340.
17. Arellano RO, Garay E, Vázquez-Cuevas FG: **Functional interaction between native G protein-coupled purinergic receptors in *Xenopus* follicles.** *Proc Natl Acad Sci USA* 2009, **106**(39):16680-16685.
18. Vázquez-Cuevas FG, Juárez B, Garay E, Arellano RO: **ATP-induced apoptotic cell death in porcine ovarian theca cells through P2X<sub>7</sub> receptor activation.** *Mol Reprod Dev* 2006, **73**(6):745-755.
19. Duleba A, Spaczynski RZ, Olive DL, Behrman HR: **Effects of insulin and insulin-like growth factors on proliferation of rat ovarian theca-interstitial cells.** *Biol Reprod* 1997, **56**:891-897.
20. Magoffin DA, Erickson GF: **Purification of ovarian theca-interstitial cells by density gradient centrifugation.** *Endocrinology* 1988, **122**(5):2345-2347.
21. Chomezynski P, Sacchi N: **Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction.** *Ann Biochem* 1987, **162**:156-159.
22. Towns R, Azhar S, Peegel H, Menon KM: **LH/hCG-stimulated androgen production and selective HDL-cholesterol transport are inhibited by a dominant-negative CREB construct in primary cultures of rat theca-interstitial cells.** *Endocrine* 2005, **27**(3):269-277.
23. Ulloa-Aguirre A, Zariñán T, Pasapera AM, Casas-González P, Dias JA: **Multiple facets of follicle-stimulating hormone receptor function.** *Endocrine* 2007, **32**(3):251-263.
24. Flores RV, Hernández-Pérez MG, Aquino E, Garrad RC, Weisman GA, Gonzalez FA: **Agonist-induced phosphorylation and desensitization of the P2Y2 nucleotide receptor.** *Mol Cell Biochem* 2005, **280**(1-2):35-45.
25. Grbic DM, Degagné E, Langlois C, Dupuis AA, Gendron FP: **Intestinal inflammation increases the expression of the P2Y6 receptor on epithelial cells and the release of CXC chemokine ligand 8 by UDP.** *J Immunol* 2008, **180**(4):2659-2668.
26. Ratchford AM, Baker OJ, Camden JM, Rikka S, Petris MJ, Seye CI, Erb L, Weisman GA: **P2Y2 nucleotide receptors mediate metalloprotease-**

- dependent phosphorylation of epidermal growth factor receptor and ErbB3 in human salivary gland cells. *J Biol Chem* 2010, **285**(10):7545-55.
27. Pierce KL, Luttrell LM, Lefkowitz RJ: **New mechanisms in heptahelical receptor signaling to mitogen activated protein kinase cascades.** *Oncogene* 2001, **20**(13):1532-1539.
  28. Schulte G, Fredholm BB: **Signalling from adenosine receptors to mitogen-activated protein kinases.** *Cell Signal* 2003, **15**(9):813-827.
  29. García-Sainz JA, Alcántara-Hernández R, Vázquez-Prado J: **α1-Adrenoceptor subtype activation increases proto-oncogene mRNA levels. Role of protein kinase C.** *Eur J Pharm* 1998, **342**:311-317.
  30. Bahr J, Kao L, Nalbandov V: **The role of catecholamines and nerves in ovulation.** *Biol Reprod* 1974, **10**:273-290.
  31. Greiner M, Paredes A, Rey-Ares V, Saller S, Mayerhofer A, Lara HE: **Catecholamine uptake, storage, and regulated release by ovarian granulosa cells.** *Endocrinology* 2008, **149**(10):4988-4996.
  32. Lara HE, Dorfman M, Venegas M, Luza SM, Luna SL, Mayerhofer A, Guimarães MA, Rosa E Silva AA, Ramírez VD: **Changes in sympathetic nerve activity of the mammalian ovary during a normal estrous cycle and in polycystic ovary syndrome: studies on norepinephrine release.** *Microsc Res Tech* 2002, **59**:495-502.
  33. Mayerhofer A, Disson GA, Costa ME, Ojeda SR: **A role for neurotransmitters in early follicular development: induction of functional follicle-stimulating hormone receptors in newly formed follicles of the rat ovary.** *Endocrinology* 1997, **138**:3320-3329.
  34. Mayerhofer A, Fritz S: **Ovarian acetylcholine and muscarinic receptors: hints of a novel intrinsic ovarian regulatory system.** *Microsc Res Tech* 2002, **59**(6):503-508.
  35. Tu MT, Luo SF, Wang CC, Chien CS, Chiu CT, Lin CC, Yang CM: **P2Y2 receptor-mediated proliferation of C6 glioma cells via activation of Ras/Raf/MEK/MAPK pathway.** *British J Pharmacol* 2000, **129**:1481-1489.
  36. Heo JS, Han HJ: **ATP stimulates mouse embryonic stem cell proliferation via protein kinase C, phosphatidylinositol 3-kinase/Akt, and mitogen-activated protein kinase signaling pathways.** *Stem Cells* 2006, **24**(12):2637-2648.
  37. Chambard JC, Lefloch R, Pouységur J, Lenormand P: **ERK implication in cell cycle regulation.** *Biochim Biophys Acta* 2007, **1773**(8):1299-1310.
  38. Salvetti NR, Panzani CG, Gimeno EJ, Neme LG, Alfaro NS, Ortega HH: **An imbalance between apoptosis and proliferation contributes to follicular persistence in polycystic ovaries in rats.** *Reprod Biol Endocrinol* 2009, **7**:68-77.
  39. Priyanka S, Medhamurthy R: **Characterization of cAMP/PKA/CREB signaling cascade in the bonnet monkey corpus luteum: expressions of inhibin-alpha and StAR during different functional status.** *Mol Hum Reprod* 2007, **13**(6):381-390.
  40. Magoffin DA: **Ovarian theca cell.** *Int J Biochem Cell Biol* 2005, **37**(7):1344-1349.
  41. Armstrong DG, Webb R: **Ovarian follicular dominance: the role of intraovarian growth factors and novel proteins.** *Rev Reprod* 1997, **2**:139-146.
  42. Coutinho-Silva R, Ojcius DM, Górecki DC, Persechini PM, Bisaggio RC, Mendes AN, Marks J, Burnstock G, Dunn PM: **Multiple P2X and P2Y receptor subtypes in mouse J774, spleen and peritoneal macrophages.** *Biochem Pharmacol* 2005, **69**(4):641-655.
  43. da Cruz CM, Ventura AL, Schachter J, Costa-Junior HM, da Silva Souza HA, Gomes FR, Coutinho-Silva R, Ojcius DM, Persechini PM: **Activation of ERK1/2 by extracellular nucleotides in macrophages is mediated by multiple P2 receptors independently of P2X7-associated pore or channel formation.** *Br J Pharmacol* 2006, **147**(3):324-334.

doi: 10.1186/1477-7827-8-88

**Cite this article as:** Vázquez-Cuevas *et al.*, Functional expression and intracellular signaling of UTP-sensitive P2Y receptors in theca-interstitial cells *Reproductive Biology and Endocrinology* 2010, **8**:88

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

