

RESEARCH

Open Access



Feedback inhibition of CREB signaling by p38 MAPK contributes to the negative regulation of steroidogenesis

Jiaxin Li^{1†}, Qian Zhou^{1†}, Zhuang Ma¹, Meina Wang¹, Wen-Jun Shen^{2,3}, Salman Azhar^{2,3}, Zhigang Guo¹ and Zhigang Hu^{1*}

Abstract

Background: Steroidogenesis is a complex, multi-steps biological process in which, cholesterol precursor is converted to steroids in a tissue specific and tropic hormone dependent manner. Given that steroidogenesis is achieved by coordinated functioning of multiple tissue specific enzymes, many steroids intermediates/metabolites are generated during this process. Both the steroid products as well as major lipoprotein cholesterol donor, high-density lipoprotein 3 (hHDL₃) have the potential to negatively regulate steroidogenesis via increased oxidative stress/reactive oxygen species (ROS) generation.

Methods: In the current study, we examined the effects of treatment of a mouse model of steroidogenesis, Y1-BS1 adrenocortical tumor cells with pregnenolone, 22(R)-Hydroxycholesterol [22(R)-diol] or hHDL₃ on ROS production, phosphorylation status of p38 MAPK and cAMP response element-binding protein (CREB), CREB transcriptional activity and mRNA expression of StAR, CPY11A1/P450scc and antioxidant enzymes, superoxide dismutases [Cu,ZnSOD (SOD1), MnSOD (SOD2)], catalase (CAT) and glutathione peroxidase 1 (GPX1). We also detected the steroid product in p38 MAPK inhibitor treated Y1 cells by HPLC-MS / MS.

Results: Treatment of Y1 cells with H₂O₂ greatly enhanced the phosphorylation of both p38 MAPK and CREB protein. Likewise, treatment of cells with pregnenolone, 22(R) diol or hHDL₃ increased ROS production measured with the oxidation-sensitive fluorescent probe 2',7'-Dichlorofluorescein diacetate (DCFH-DA). Under identical experimental conditions, treatment of cells with these agents also increased the phosphorylation of p38 MAPK and CREB. This increased CREB phosphorylation however, was associated with its decreased transcriptional activity. The stimulatory effects of pregnenolone, 22(R)-diol and hHDL₃ on CREB phosphorylation was abolished by a specific p38 MAPK inhibitor, SB203580. Pregnenolone, and 22(R) diol but not hHDL₃ upregulated the mRNA expression of SOD1, SOD2 and GPX1, while down-regulated the mRNA levels of StAR and CYP11A1. The p38 inhibitor SB203580 could increase the steroid production in HDL₃, 22(R)-diol or pregnenolone treated cells.

Conclusion: Our data demonstrate induction of a ROS/p38 MAPK-mediated feedback inhibitory pathway by oxy-cholesterol and steroid intermediates and products attenuates steroidogenesis via inhibition of CREB transcriptional activity.

Keywords: Steroidogenesis, p38 MAPK, CREB, Steroids intermediates, Feedback regulation

* Correspondence: huzg_2000@126.com

†Equal contributors

¹Jiangsu Key Laboratory for Molecular and Medical Biotechnology, College of Life Sciences, Nanjing Normal University, 1 WenYuan Road, Nanjing 210023, China

Full list of author information is available at the end of the article



Background

Steroidogenesis is a multi-step process by which the cholesterol is converted to parent steroid, pregnenolone, which is further metabolized into other steroids in a tissue specific manner [1, 2]. The cholesterol required for steroid hormone synthesis can be theoretically obtained from several different potential sources including de novo synthesis from acetate, cholesteryl esters stored in the form of lipid droplets or can be obtained from circulating lipoproteins via low-density lipoprotein (LDL) receptor/endocytic pathway or SR-BI (for high-density lipoprotein or HDL)/selective uptake pathway [3, 4]. Steroidogenic process is subjected to a dual regulation—acute and chronic regulation [5], although both are under the control of tissue-specific tropic hormone. Adrenocorticotropin hormone (ACTH) increases glucocorticoid (cortisol in humans and corticosterone in rodents) synthesis in adrenal fasciculata cells, ACTH, K^+ or angiotensin II (AngII) control mineralocorticoid (aldosterone) synthesis in adrenal glomerulosa cells, follicle-stimulating hormone (FSH) controls the female sex steroid (progesterone and estrogen) synthesis in ovarian granulosa cells, whereas luteinizing hormone (LH) regulates progesterone synthesis in luteinized ovarian granulosa-luteal cells, androgen production in ovarian theca-interstitial cells and testosterone synthesis in testicular Leydig cells [2].

ROS, such as H_2O_2 and superoxide anion ($O_2^{\bullet-}$) are produced by the mitochondrial electron transport chain as a byproduct of oxidative phosphorylation [6]. In steroidogenic cells, there is a secondary source for ROS production. During steroidogenesis, ROS is produced by cytochrome P450 enzymes catalyzing the steroid hydroxylation steps, particularly by CYP11B1 and to a lesser extent by CYP11A1 [7]. Moreover, tropic hormones, ACTH and LH not only stimulate steroid synthesis in steroidogenic cells of their target tissues but also promote ROS production, which in turn can cause DNA damage, protein oxidation and membrane lipid peroxidation in steroidogenic cells [8, 9]. We have previously reported that age-related decline in steroid synthesis is caused by excessive oxidative damage to cellular machinery involved in steroidogenesis. In addition, we demonstrated that aging leads to a significant reduction in enzymatic and non-enzymatic antioxidant systems and increased membrane lipid peroxidation in steroidogenic cells of adrenal gland and testis [10].

Extensive evidence now suggests that low (physiological) but not high concentrations of ROS such as H_2O_2 contribute to the regulation of multiple cellular signaling pathways [11–13]. Given this, we considered the possibility that ROS generated during steroidogenesis may have a negative feedback effect on steroid hormone synthesis. Indeed, there is evidence that ROS such as H_2O_2 interferes with the normal transport of

mobilized cytoplasmic cholesterol to and within the mitochondria for side chain cleavage in steroidogenic cells [14–16]. Current evidence suggests that one mechanism by which ROS may influence cellular signaling pathways, is via increased phosphorylation and activation of oxidant sensitive p38 MAPKs [17–19]. p38 MAPKs are activated by a wide variety of cellular stresses, such as cytokines, ultraviolet irradiation, heat shock, and osmotic shock, and in turn regulate a number of metabolic processes including cell proliferation, differentiation, apoptosis and other numbers of different biological effects [20–22]. Our own studies have shown that oxidant mediated activation of p38 MAPK inhibits steroidogenesis via down-regulation of StAR gene transcription [22, 23].

Besides oxidants, there is ample evidence to suggest that tropic hormones (e.g., FSH and ACTH) themselves can stimulate the phosphorylation and activation of p38 MAPK. The hormonal activation of p38 MAPK, is presumably required to terminate the hormone-stimulated acute steroidogenesis. When considered in this context, it is likely that p38 MAPK serves as a negative regulator of steroidogenesis via a feedback dependent mechanism. In the current study, we sought to determine whether the steroid intermediates or precursors can stimulate p38 MAPK phosphorylation/activation. We provide evidence that the steroid substrate/precursors and intermediates, HDL₃, 22(R)-diol and pregnenolone, can promote the phosphorylation and activation of p38 MAPK. In addition, phosphorylation and activity of CREB protein, a key transcription factor involved in the transcriptional regulation of steroidogenic enzymes, is also regulated by these metabolites. The mRNA expression of StAR and CYP11A1 (P450scc), which mediate the first steps in steroidogenesis, were detected and regulated by these steroid precursors and intermediates. Finally, we demonstrate that expression of antioxidant enzymes, SODs and GPX is also subjected to regulation by the steroid substrate/precursors and intermediates. In summary, this study provides first evidence that treatment of a model rodent adrenal cell line with steroid precursors/ metabolites promote ROS production, stimulate (activate) p38 MAPK and CREB phosphorylation and changed gene transcription of StAR, CYP11A1 and three major antioxidant enzymes. The ROS/p38 MAPK-mediated increased phosphorylation of CREB, however, is associated with a reduction in its transcriptional activity. This attenuation in CREB transcriptional activity in turn results in inhibition of mRNA expression of key steroidogenic enzymes/proteins such as StAR via a feedback mechanism.

Methods

Reagents and antibodies

22(R)-Hydroxycholesterol [22(R)-diol], 21-Acetoxyprogesterone and p38 MAP kinase inhibitor (SB 203580)

were obtained from Sigma–Aldrich (St. Louis, MO). Promega Dual-Luciferase Reporter Kit (E1980) was purchased from Promega Corporation (Madison, WI). pCRE-Luc cis-reporter plasmid and pLuc-MCS vector were obtained from Agilent Technologies (Santa Clara, CA, USA). 20 α -hydroxyprogesterone was purchased from Chemsky International Co., Ltd. (Shanghai, China). Anti-phospho-CREB (Ser133), anti-CREB, anti-p38 MAPK, and anti-phospho-p38 MAPK (Thr180/Tyr182) antibodies were supplied by Cell Signaling Technology, Inc. (Danvers, MA, USA). Anti-GAPDH was purchased from Santa Cruz Biotechnology (Santa Cruz, CA, USA). IRDye[®] 800CW Goat anti-Mouse IgG (H + L) and IRDye[®] 800CW Goat anti-rabbit IgG (H + L) were purchased from LI-COR Biosciences (Lincoln, NE, USA). SYBR Green Master Mix was supplied by TAKARA Biotechnology (Dalian) Co. LTD (Dalian, China). Lipofectamine[®] 2000 was obtained from Invitrogen (Life Technologies, Grand Island, NY, USA). Tissue culture supplies were from Life Technologies through its Gibco Cell Culture Media Division (Grand Island, NY, USA).

Cell culture and treatment

SR-BI enriched Y1-BS1 cells were initially kindly supplied by late Dr David Williams (State University at Stony Brook, Stony Brook, NY, USA). Y1-BS1 cells, which is response to hormone and lipoprotein with increased steroidogenesis, is a stable subclone of the Y1 mouse adrenocortical tumor cell line isolated by Yasumura et al. 1966 [24, 25]. Y1-BS1 has been proven to be robust in steroidogenesis and retains responsiveness to hormone [24]. In contrast to mouse adrenocortical cells/adrenal gland that produce corticosterone as the major steroid product, cultured Y1-BS1 cells, both under basal conditions and in response to trophic hormone stimulation, synthesize and secrete 20 α -hydroxy- Δ 4-pregnene-3-one (20 α -hydroxyprogesterone) and 11 β ,20 α -dihydroxy- Δ 4-pregnene-3-one (11 β , 20 α -hydroxyprogesterone) [26]. Y1-BS1 cells also secrete small quantities of progesterone. Y1-BS1 cells were cultured in F-12K medium supplemented with 10% fetal bovine serum (FBS), and 100 unit/ml penicillin and 100 μ g/ml streptomycin. Cell cultures were maintained at 37 °C in a humidified incubator in the presence of 5% CO₂/95% air. When required, cells were incubated with H₂O₂ (100 nM) or Bt2cAMP (2.5 mM) for an appropriate time [27]. For other treatment, cells were plated and cultured in 6-well or 12-well plates with 10% bovine (b) lipoprotein-deficient serum (LPDS), and subsequently treated with human (h) apoE-free high density lipoproteins (hHDL₃, 30 μ g/ml), 22(R)-diol (10 μ M) or pregnenolone (10 μ M) for varying incubation time. hHDL₃ and bLPDS were isolated as previously described [28, 29].

Measurement of intracellular reactive oxygen species (ROS) production

The levels of intracellular ROS were quantified using the Reactive Oxygen Species Assay Kit purchased from Beyotime Biotechnology (Shanghai, China). In brief, triplicate dishes of cultured Y1-BS1 cells (5X10⁵ cells/well) were pre-loaded with 10 μ M DCFH-DA (2,7-Dichlorodihydrofluorescein diacetate) for 30 min. Subsequently, dishes were washed and incubated with HDL₃ (30 μ g protein/ml), 22(R)-diol (10 μ M) or pregnenolone (10 μ M) for indicated time. At the end of incubation, the culture medium was removed from dishes and cells were immediately lysed and centrifuged. DCFH-DA fluorescence of the cell lysates was measured using Tecan Infinite[®] 200 Pro Microplate Reader with excitation and emission wavelengths of 480 and 525 nm respectively.

Luciferase assays

Luciferase assays were carried out using cell extracts from transfected Y1-BS1 cells. Groups of cultured dishes with 60% confluent Y1-BS1 cells were transfected with 1 μ g pCRE-luc plasmid or pLuc-MCS vector per well in a 24-well plate using Lipofectamine[®] 2000 as a transfection reagent (Life Technologies, Grand Island, NY, USA). Twenty-four hours after transfection, the cells were re-cultured in F-12K medium supplemented with 10% LPDS overnight followed by treatment with HDL₃, 22(R)-diol or pregnenolone for an additional 5 h. Cells were harvested, lysed and cell extracts luciferase assayed with dual-luciferase reporter assay system (Promega) according to the manufacturer's protocol and light signal (bioluminescence) was quantified using a Tecan Infinite[®] 200 Pro luminometer. Renilla luciferase was used as a normalization control. The results are expressed as relative luciferase activity (ratio of Firefly/Renilla luciferase activity), and data shown are the mean (\pm SD) of triplicate values obtained from a representative experiment that was independently repeated for at least three times.

RNA isolation and quantitative PCR

Total RNA was extracted from Y1-BS1 cells using a miRNeasy mini kit (Qiagen Inc., Valencia, CA) following manufacturer's instructions. The isolated RNA was transcribed to first strand cDNA at 42 °C for 1 h in an incubation medium containing 2 μ g of total RNA and superscript II reverse transcriptase (Life Technologies, Grand Island, NY). Amplification of cDNAs was performed with an ABI StepOneplus system according to manufacturer's instructions. Each sample contained 2 μ l cDNA (1:10 dilution of the original cDNA), 500 nM each of sense and antisense primer, 10 μ l 2x SYBER Green premix and 0.4 μ l Rox (qPCR kit, cat. no.

RR420A; TAKARA Biotechnology, Dalian, China) in a final volume of 20 μ l. 36B4 was used as an internal control. All primer sequences used for qPCR are presented in Table 1.

Western blot analysis

For Western blotting, Y1-BS1 cell lysates were mixed with equal volumes of 5 \times Laemmli sample buffer [120 mM Tris-HCl, pH 6.8, 2% SDS (w/v), 10% sucrose (w/v), and 1% 2-mercaptoethanol] and subjected to 12% SDS-PAGE. For each sample, a constant amount of protein (10–20 μ g) was loaded on the gel. Protein markers were also loaded on the gel. After electrophoretic separation, the proteins were transferred to Immobilon PVDF membranes (Millipore Corp., Bedford, MA) using standard techniques. The protein blots were incubated with first antibody for 2 h at room temperature, then probed with IRDye infrared secondary mouse or rabbit anti-rabbit IgG and visualized using the odyssey[®] infrared imaging system (LI-COR Biosciences, Lincoln, NE, USA).

Hormone analysis

Y1-BS1 cells were pre-loaded into 12-well plate (5 \times 10⁵ cells/well) and then cultured in F-12K medium supplemented with 10% LPDS \pm p38 MAPK inhibitor SB203580 overnight. Subsequently, dishes were washed and incubated with hHDL₃ (30 μ g protein/ml), 22(R)-diol (10 μ M) or pregnenolone (10 μ M) for 5 h. The medium samples were collected for hormone analysis using isotope dilution high performance liquid chromatography-tandem mass spectrometry (ID-HPLC-MS / MS). HPLC-MS / MS was performed by Shanghai Dian Medical Testing Institute (Shanghai, China) as previously described [30].

Table 1 Primers used for quantitative real-time PCR

Primers for qPCR	
Mouse StAR	5'-CGGAGCAGAGTGGTGCATC-3'-F 5'-TGAGTTTAGTCTTGGAGGGACTTC-3'-R
Mouse CYP11A1	5'-ACTGTGAACCTGAAGCTGG-3'-F 5'-GGGAAAGAGGGAAAGAGGATG-3'-R
Mouse SOD1	5'-AAGACTGGAAATGCTGGGAG-3'-F 5'-GGTTTGAGGGTAGCAGATGAG-3'-R
Mouse SOD2	5'-TGCTCTAATCAGGACCCATTG-3'-F 5'-CATTCTCCAGTTGATTACATTCC-3'-R
Mouse CAT	5'-TCACCTGTAATCAACGCTGG-3'-F 5'-AGCCCTAACCTTTCATTTCCC-3'-R
Mouse GPX1	5'-CAGGAGAATGGCAAGATGAAG-3'-F 5'-GAAGGTAAGAGCGGGTGAG-3'-R
Mouse 36B4	5'-TTTGGGCATCACCACGAAAA-3'-F 5'-GGACACCTCCAGAAAGCGA-3'-R

Statistical analysis

Data are expressed as mean \pm SEM for at least three independent experiments. Statistical analyses were performed using ANOVA followed by the Bonferroni's post-test using GraphPad Prism Software, Prism 6 (GraphPad Software, Inc., San Diego, CA, USA). A statistical difference of $p < 0.05$ was considered significant.


Results

H₂O₂ stimulates the phosphorylation of p38 MAPK and cAMP responsive CREB transcription factor

Increasing evidence suggests that ROS such as \bullet O₂⁻, \bullet OH and H₂O₂, promote the phosphorylation and activation of MAPK, among which p38 MAPK is highly responsive to oxidant stress which has important roles in cell signaling and homeostasis [31–34]. We previously demonstrated that treatment of Y1-BS1 cells with superoxide, H₂O₂ or a lipid peroxidation product, 4-hydroxy-2-nonenal (HNE) reciprocally inhibited steroid production and increased the phosphorylation and activation of p38 MAPK [23]. CREB is a key transcription factor that plays a pivotal role in the regulation of steroidogenic enzymes at their gene transcription level. It is stimulated by a number of agents including cAMP, growth factors and UV exposure [35–37]. The results presented in Fig. 1a and b demonstrate that treatment of Y1-BS1 cells with 100 nM H₂O₂ for 5 min caused a robust phosphorylation of p38 MAPK and a modest but significant stimulation of phosphorylated form of CREB for 20 min, thereafter the phosphorylation of p38 and CREB decreased. Pretreatment of cells with a specific p38 MAPK inhibitor, SB203580 abolished the stimulatory effects of H₂O₂ on p38 MAPK and CREB phosphorylation. These results not only complement the previously published studies from our laboratory [22] but also unequivocally establish that CREB is a downstream critical target of p38 MAPK in steroidogenic cells. Meanwhile, we detected that Bt₂cAMP could stimulate both the phosphorylation of p38 MAPK and CREB (Fig. 1c). While SB203580 abolished the stimulatory phosphorylation of p38 MAPK by Bt₂cAMP, phosphorylation of CREB was not diminished (Fig. 1d). These results are consistent with the reported finding that cAMP stimulated the phosphorylation of CREB through PKA pathway [36, 38].

Steroid intermediates, 22(R)-diol and pregnenolone and circulating cholesterol carrier hHDL₃ promote intracellular ROS production

Steroidogenesis being a multi-step process results in generation and/or accumulation of a number of intermediary metabolites such as 22(R)-diol and pregnenolone, which could potentially further enhance the intracellular ROS production. This possibility is in line




with the earlier evidence showing that the steroid intermediates, testosterone, testosterone precursors or analogs e.g., epitestosterone and 17 α -methyltestosterone, can cause oxygen tension sensitive decrement in P450 hydroxylase activity [39]. In addition, the cholesterol carrier, hHDL₃ may also directly or indirectly influence the ROS production, especially since HDL is known to participate in cellular signaling [40–42] and is the predominant form of circulating lipoprotein in rodent and a major source of cholesterol for steroidogenesis. Here, we tested the ability of HDL₃, 22(R)-diol and pregnenolone to modulate ROS production in Y1-BS1 cells. For these studies, Y1-BS1 cells were cultured in regular F-12K medium containing 10% FBS for a brief period of time to allow the cells to attach to the plate and then culture medium was replaced with a fresh medium containing 10% LPDS and dishes incubated overnight. Subsequently, the cells were treated ± HDL₃, ±22(R)-diol or ± pregnenolone in the presence of 10% LPDS for various time points. At the end of incubation, intracellular ROS generation was evaluated using DCFH-DA assay. The results depicted in Fig. 2 demonstrate that HDL₃, 22(R)-diol or and pregnenolone, all stimulated ROS generation

with a biphasic response; an initial 10 min rapid response and a gradually declining slow phase during the next 60 min (Fig. 2).

Treatment of Y1-BS1 cells with steroid linked intermediates/metabolites and hHDL₃ results in increased phosphorylation of p38 MAPK

We next tested the ability of HDL₃, 22(R)-diol and pregnenolone to promote the phosphorylation of p38 MAPK. Treatment of Y1-BS1 with hHDL₃ (30 μ g protein/ml), 22(R)-diol (10 μ M) or pregnenolone (10 μ M) increased p38 MAPK phosphorylation in a time dependent manner (Fig. 3a). In each case, the maximum stimulatory effect of the agent was noted between 20 and 40 min after exposure and thereafter phosphorylation levels of p38 MAPK began to decline reaching to the basal levels at the end of 240 min of incubation. The total p38 MAPK levels, however, were not impacted by treatment of the cells with any of the three agents (Fig. 3a). We next evaluated whether hHDL₃ and/or steroid metabolites can modulate CREB phosphorylation via p38 MAPK. CREB is one of the best understood phosphorylation dependent transcription factors [43–45]. It



activates transcription of the target genes in response to a diverse array of stimuli including peptide hormones, growth factors and neuronal activity. Figure 3b shows that HDL₃, 22(R)-diol or pregnenolone treatment, like H₂O₂, increased the phosphorylation of CREB with maximal stimulation occurring around 20 min. These results led us to conclude that HDL₃, 22(R)-diol and pregnenolone promote CREB phosphorylation via upstream p38 MAPK signaling pathway. To provide additional evidence for this conclusion, we treated cells with these three agents in the presence and absence of a specific p38 MAPK inhibitor, SB203580 and reassessed the phosphorylation status of both p38 MAPK and CREB proteins. The results presented in Fig. 4 demonstrate that co-treatment of cells with SB203580 blocked the ability of HDL₃, 22(R)-diol and pregnenolone to stimulate the phosphorylation of both p38 MAPK and CREB, further confirming the p38 MAPK catalyzed phosphorylation of CREB.

Steroid intermediates repress CREB activity

The above results indicate that phosphorylated forms of p38 MAPK and CREB were significantly increased in Y1-BS1 exposed to HDL₃, 22(R)-diol or pregnenolone compared to vehicle control (Figs. 3 and 4). We next evaluated the impact of these agents on CREB activity using a CRE/CREB Reporter kit designed for monitoring the activity of the cAMP/PKA signaling in cultured cells. As shown in Fig. 5, CREB activity was slightly but significantly decreased in response to treatment with these three agents for 3 to 5 h. These data open up the

possibility that overproduction/increased accumulation of steroid intermediates or sustained activation of cells to HDL leads to enhanced ROS production and consequently loss of CREB activity.

Regulation of StAR and CYP11A1 gene expression

StAR protein mediates rate-limiting step of steroidogenesis in which cholesterol was translocated from cytoplasm to mitochondria [46]. Then, P450_{scc} (CYP11A1) catalyzes the conversion of StAR-delivered cholesterol to pregnenolone [47]. The transcription of both StAR and CYP11A1 is also regulated by tropic hormones involving cAMP signaling pathway in multiple transcription factors, including CREB [48]. Since we observed that CREB activity is repressed by HDL₃, 22(R)-diol and pregnenolone, we measured the mRNA levels of its downstream targets, StAR and CYP11A1 following treatment of cells with these three agents. As shown in Fig. 6, the mRNA expression of both StAR and CYP11A1 was down-regulated by 22(R)-diol and pregnenolone treatment but not by hHDL₃ for 1 or 3 h' treatment, while the mRNA expression recovered after a longer time treatment.

Inhibition of p38 MAPK phosphorylation upregulate steroidogenesis

We have demonstrated that inhibition of p38 MAPK activity with SB203580 potentiated the Bt₂cAMP and Bt₂cAMP + hHDL₃-stimulated steroid production in Y1-BS1 adrenal cells [23]. Although, oxidants-induced activation p38 MAPK activity critically mediates the oxidant inhibition of steroid production, we also evaluated the potential effects of steroid metabolites and intermediates -induced activation of p38 MAPK on steroid synthesis. Y1-BS1 cells were treated with three steroid intermediates in the presence and absence of SB203580 and medium samples analyzed for steroid production using HPLC-MS/MS. As shown in Fig. 7, although additional substrates (HDL₃, 22(R)-diol or pregnenolone) loading increased the steroid production, inclusion of SB203580 to inhibit the p38 activity could increase the steroid production in HDL₃, 22(R)-diol or pregnenolone treated cells. These findings are consistent with previous results showing that inhibition of p38 MAPK by SB203580 could ameliorate the oxidative stress-induced repression of steroidogenesis [23].

Upregulation of anti-oxidant enzymes gene expression

ROS-induced oxidative stress is defined as a disturbance in the balance between production of ROS and antioxidants defenses, particularly the reduced functioning of the antioxidant enzymes (e.g., SOD1 [cytosolic CuZn-SOD], SOD2 [mitochondrial Mn-SOD], GPX1 [cytosolic/mitochondrial glutathione peroxidase 1], CAT [peroxisomal catalase] and PRDX3 [mitochondrial




Fig. 3 Steroid intermediates promote phosphorylation of p38 MAPK (a) and CREB (b) in a time dependent manner. Y1-B51 cells were incubated with HDL₃ (30 μg/ml), 22(R)-diol (10 μM) or pregnenolone (10 μM) for different time point. At the end of incubation, cells were harvested and lysed for SDS-PAGE followed by western blotting analysis using Phospo-specific and total p38 MAPK and CREB antibodies. Data shown are representative of three independent experiments

peroxiredoxin 3] [7, 8, 49–52]. Here we evaluated the mRNA expression levels of SOD1, SOD2, CAT and GPX1 expression in Y1-B51 cells following their treatment with HDL₃, 22(R)-diol or pregnenolone. Contrary to our expectation, we were surprised to see that 22(R)-diol and pregnenolone but not hHDL₃ treatment induced the mRNA expression of SOD1, SOD2 and GPX (Fig. 8). In contrast, catalase expression was not affected by any of these treatments.

Discussion

Steroidogenesis represents a highly complex and multi-step metabolic process involving the participation of several catalytic enzymes, regulatory molecules and cellular organelles. More specifically, it involves the participation of various steroidogenic enzymes (e.g., CYP11A1, CYP11B1, 3βHSD), cholesterol transport proteins (e.g., SNAREs, StAR, vimentin), transcription factors (SF-1, DAX-1, AP-1, CREB and others) and availability of adequate amounts of substrate cholesterol [1, 2, 48, 53, 54]. Although tissue specific tropic hormones are the principal

regulators of steroidogenesis, many growth factors, cytokines and pathophysiological conditions such as oxidative stress and ER stress also contribute to the regulation of steroidogenesis [1, 15, 53]. Because steroid synthesis involves many enzymatic steps, a number of intermediate steroid hormone metabolites are generated and sequentially utilized for the production of tissue specific steroid hormone. However, for many of these intermediates, conversion to subsequent metabolite is not efficient resulting in their increased accumulation in steroidogenic cells and as such could potentially impact steroidogenesis via increased ROS formation. Besides, steroidogenesis itself significantly contributes to the mitochondrial ROS production and in turn excessive oxidative stress can impede steroidogenesis [7, 8, 10, 23, 55]. As noted before, ROS are derived from molecular oxygen (O₂) and comprise molecules with varying oxidant properties. At low concentrations, ROS modulate many cellular processes through redox-dependent signaling including proliferation, differentiation, apoptosis, immune regulation, cellular adaptation [11–13] and now it is becoming clear that they also






Fig. 4 Inhibition of p38 MAPK blocked the phosphorylation of CREB by steroid intermediates. Y1-BS1 cells were pre-treated without (a) or with (b) p38 MAPK inhibitor (10 μ M) overnight. Cells were then incubated with HDL₃ (30 μ g/ml), 22(R)-diol (10 μ M) or pregnenolone (10 μ M) for 20 min. At the end of the incubation, cell lysates were subjected to SDS-PAGE followed by western blotting analysis. Data shown are representative of three independent experiments

participate in the (negative) regulation of steroidogenesis [7, 8, 10, 23, 55, 56]. In contrast, overproduction or high levels of ROS results in oxidative stress, a deleterious process that can be an important mediator of damage to cellular macromolecular structures including lipids and membranes, proteins and DNA [57, 58]. Thus, mammalian cells and more specifically, steroidogenic cells possess a potent anti-oxidant network to scavenge and neutralize the excessively produced ROS. Under normal physiological conditions, a balance between ROS production and elimination maintains optimal homeostasis. However, disturbances of such equilibrium under pathological conditions can alter the functioning of normal cellular process with detrimental metabolic consequences.

In this study we made a novel observation that endogenous steroid intermediates such as 22(R)-diol and pregnenolone as well as a donor of cholesterol substrate, hHDL₃ can promote ROS production and ensuing oxidative stress in Y1-BS1 adrenal cells. Our observations are in line with previous studies showing superoxide/ H₂O₂ produced during ACTH-stimulated adrenal steroid production is involved in the termination of the steroidogenic response [8]. They demonstrated that H₂O₂ produced by P450 enzymes during steroidogenesis inactivates potent mitochondrial antioxidant enzyme, peroxiredoxin 3, which in turn triggers a sequence of events including increased accumulation of H₂O₂,

activation of p38 MAPK, suppression of StAR mRNA/protein expression and inhibition of steroidogenesis [8]. Our results suggest that it is not only the H₂O₂ produced during Mn-SOD catalyzed dismutation of superoxide anions and ensuing inactivation of peroxiredoxin 3 that inhibits steroidogenesis, but also steroid metabolites such as 22(R)-diol and pregnenolone possess similar capabilities in promoting ROS production and interfering with normal steroidogenesis. However, we were surprised to find that hHDL₃ a cholesterol donor for steroidogenesis also promotes ROS production. This is even more surprising given that HDL is considered an antioxidant with relevance to cardiovascular disease [59]. At present we are unable to provide exact explanation by which HDL treatment of Y1 BS1 cells results in increased ROS production, but it may be that hHDL₃ being a cholesterol donor, delivers excessive amounts of cholesterol to cells for its catabolism and this leads to exaggerated production of ROS and excessive oxidative stress. Obviously, more experimental studies are needed to sort out among these various possibilities.


Previous data suggest that hydrogen peroxide is a potent activator for p38 MAPK [23, 60]. Other reports indicate that tropic hormones such as FSH and ACTH also induce the phosphorylation and activation of p38 MAPK [8, 61]. Several investigators also reported that p38 MAPK is one of the regulators of CREB phosphorylation and its activity [22, 36, 62]. Our data provide




recruitment of CBP or p300 and could not activated the transcription of CREB-dependent reporters [38]. A number of studies have reported that CREB is a target of p38 MAPK and that it phosphorylates and activates the activity of CREB [36, 62, 65]. On the contrary, other studies including our own have shown that p38 MAPK inhibits the transcriptional (reporter) activity of CREB in model steroidogenic cell lines, MLTC and Y1, while at the same time promotes CREB phosphorylation [22, 36, 62]. It is likely that these variable findings about p38 MAPK-mediated phosphorylation and altered CREB activity may be due to differences in cell types, metabolic conditions of the cell under study and functional interaction of CREB with specific co-activators/co-suppressors and/or other transcription factors. Our results provide evidence confirming previous findings from the laboratory that p38 MAPK inhibits steroidogenesis by attenuating CREB activity in a classic feedback mechanism.

Currently, we are exploring the potential mechanism by which ROS/p38 MAPK inhibition of CREB activity may lead to impaired steroidogenesis. One possibility we are considering is the inhibition of StAR gene transcription as a result of oxidant-mediated impaired CREB

evidence that HDL₃, 22(R)-diol or pregnenolone treatment of Y1-BS1 results in robust phosphorylation of p38 MAPK as a consequence of enhanced ROS production. Under identical experimental conditions, the phosphorylation of one of the important steroidogenic transcription factors, CREB was also induced and based on the use of inhibitors, the observed phosphorylation of CREB is mediated by p38 MAPK. This observation is in agreement with previous reports showing that ROS upregulates CREB phosphorylation via p38 MAPK signaling cascade [22, 63]. Although cAMP-activated PKA and angiotensin II (AngII) activated protein kinase D (PKD) catalyzed phosphorylation of CREB at Ser133 is essential for CREB-induced gene transcription, the p38 MAPK-dependant phosphorylation of CREB have yielded conflicting results [48, 64]. Naqvi et al. reported that PKA catalyzed CREB phosphorylation and promoted the recruitment of the co-activator proteins CBP (CREB-binding protein) and p300 to increase the transcription of CREB-dependent genes in MEF cells, while MSK/MAPK also phosphorylated CREB without promoting



activity. This is in line with the observation that StAR expression is sensitive to both physiological and pathophysiological levels of ROS. Moreover, earlier studies have shown that CREB protein in cooperation with SF1 is a major regulator of StAR protein gene transcription [66, 67]. Finally, we have earlier shown that oxidants-p38 MAPK cause inhibition of StAR promoter activity primarily by interfering with CREB activity [22]. Other investigators reported that HDL₂, very-low-density lipoprotein (VLDL) and glyco-oxidized VLDL can induce Cyp11B2 expression and stimulate steroid production in a human adrenocortical carcinoma cell line, NCI H295R [68–70]. Saha et al [70] also reported slight increases in StAR expression by native VLDL and glycol-oxidized VLDL but not by oxidized VLDL. The data presented here show that 22(R)-diol and pregnenolone but not HDL₃ repress the gene expression of StAR and CYP11A1. We further provided the evidence that 22(R)-diol and pregnenolone-mediated repression of StAR and CYP11A1 gene expression is achieved through excessive oxidative stress and associated p38 MAPK signaling cascade. Our steroid hormone production data provide additional support to the notion that SB203580 inhibition of p38 MAPK augments the steroids production in Y1-BS1 cells treated with steroid metabolites/intermediates. A number of studies have implicated p38 MAPK signaling cascade in the regulation of steroidogenesis [71], although p38 MAPK regulation of steroidogenesis is



complex and extent of p38 MAPK varies with steroidogenic cell types. For example, inhibition of p38 MAPK activity by SB203580 in IL-1α-stimulated immature rat Leydig cells leads to downregulation of StAR gene expression and attenuation of steroid production. [72].

Other studies have shown that inhibition of p38 MAPK activity by SB203580 in ovarian granulosa cells is accompanied by increased inhibition of LH/hCG/FSH mediated StAR expression and progesterone synthesis [61, 73]. Likewise, in primary cultures of rat adrenal glomerulosa cells, Angiotensin II activates the p38 MAPK and results in increases in StAR expression and steroid synthesis [71, 74]. Interestingly, our previous studies have shown that inhibition of p38 MAPK by either SB203580 or SB202190 in adrenocortical from old rats restores corticosterone synthesis to the levels seen in cells from young animals [23]. In this study, inhibition of p38 MAPK activity by SB203580 enhanced 20 α -hydroxyprogesterone production in mouse Y1-BS1 adrenocortical tumor cell cotreated with HDL₃, 22(R)-diol or pregnenolone. In addition, we will examine whether ROS/p38 MAPK modulate the expression of some of the critical SNARE proteins. In a recent publication, we identified several SNAREs, whose expression is essential for cholesterol transport to outer mitochondrial membrane for optimal steroid production [54].

Another surprising finding from our studies was that treatment of Y1-BS1 with 22(R)-diol and pregnenolone but not HDL₃ leads to increased mRNA expression of three antioxidant enzymes, SOD1, SOD2 and CAT. In contrast, studies by Kil et al [8] observed no changes in the expression of levels of SOD1, SOD2, CAT, and GPX1 in intact adrenals following treatment of mice with ACTH. Interestingly, a recently published study reported that p38 MAPK α causes the induction of antioxidant enzymes SOD2 and catalase by two distinct mechanisms [18]. Obviously, more studies are needed to sort out molecular mechanisms involved in the transcriptional/posttranscriptional and/or post-translational regulation of these antioxidant enzymes *in vitro* and *in vivo*.

Conclusion

In conclusion, our studies provide evidence that exposure of adrenal cells to steroid intermediates/metabolites, 22(R)-diol and pregnenolone, and hHDL₃ led to increased ROS production and associated enhanced phosphorylation of p38 MAPK and CREB. This oxidant-mediated up-regulation of CREB phosphorylation is mediated by p38 MAPK. The increased CREB phosphorylation however, was accompanied by a significant loss of CREB's transcriptional activity. Furthermore, treatment of cells with 22(R)-diol and pregnenolone and ensuing oxidative stress resulted in decreased mRNA levels of StAR and CYP11A1 and increased mRNA levels of antioxidant enzymes SOD1, SOD2 and CAT. From these studies we conclude that ROS/p38 MAPK inhibition of CREB transcriptional activity is likely responsible for ROS-induced feedback inhibition of steroidogenesis.

Abbreviations

22(R)-diol: 22(R)-Hydroxycholesterol; ACTH: Adrenocorticotropin hormone; AngII: Angiotensin II; CaM II: Calcium-calmodulin kinase II; CAT: Catalase; CREB: cAMP response element-binding protein; DCFH-DA: 2',7'-Dichlorofluorescein diacetate; FBS: Fetal bovine serum; FSH: Follicle-stimulating hormone; GPX: Glutathione peroxidase; HDL: High-density lipoprotein; HNE: 4-hydroxy-2-nonenal; IMM: Inner mitochondrial membrane; LDL: Low-density lipoprotein; LH: Luteinizing hormone; LPDS: Lipoprotein-deficient serum; MSK: Mitogen- and stress-activated kinase; OMM: Outer mitochondrial membrane; PI3K: Phosphatidylinositol 3-kinase; PKA: Protein kinase A; PKB: Protein kinase B; PKC: Protein kinase C; PKD: Protein kinase D; pp90 RSK: pp 90 ribosomal S6 kinase; ROS: Reactive oxygen species; SF1: Steroidogenic factor 1; SOD: Superoxide dismutase; SR-BI: Scavenger Receptor Class B, Type I; VLDL: Very-low-density lipoprotein

Acknowledgement

None

Funding

The work is supported by National Natural Science Foundation of China (31400659), Jiangsu Provincial Natural Science Foundation (BK20140920), Priority Academic Program Development of Jiangsu Higher Education Institutions, Natural Science Fund of Colleges and Universities in Jiangsu Province (14KJB180012), and National Institute of Health (NIH/NHLBI 2R01HL033881 and VA Merit Review 5I01BX001923, SA).

Availability of data and materials

Not applicable.

Authors' contributions

Conceived and designed the experiments: JL, ZG and ZH. Performed the experiments: JL, QZ, ZM and ZH. Analyzed the data: JL, QZ, ZM, MW and ZH. Wrote the paper: ZH, SA. Edited the manuscript: ZH, WJS, ZG and SA. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable.

Ethics approval and consent to participate

Not applicable.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Jiangsu Key Laboratory for Molecular and Medical Biotechnology, College of Life Sciences, Nanjing Normal University, 1 WenYuan Road, Nanjing 210023, China. ²Geriatric Research, Education and Clinical Center, VA Palo Alto Health Care System, Palo Alto, CA 94304, USA. ³Stanford University School of Medicine, Palo Alto, CA 94304, USA.

Received: 2 December 2016 Accepted: 6 March 2017

Published online: 16 March 2017

References

1. Miller WL, Auchus RJ. The molecular biology, biochemistry, and physiology of human steroidogenesis and its disorders. *Endocr Rev.* 2010;32:81–151.
2. Hu J, Zhang Z, Shen W-J, Azhar S. Cellular cholesterol delivery, intracellular processing and utilization for biosynthesis of steroid hormones. *Nutr Metab.* 2010;7:1.
3. Azhar S, Reaven E. Scavenger receptor class BI and selective cholesteryl ester uptake: partners in the regulation of steroidogenesis. *Mol Cell Endocrinol.* 2002;195:1–26.
4. Hattangady NG, Olala LO, Bollag WB, Rainey WE. Acute and chronic regulation of aldosterone production. *Mol Cell Endocrinol.* 2012;350:151–62.

5. Lehoux J-G, Fleury A, Ducharme L. The acute and chronic effects of adrenocorticotropin on the levels of messenger ribonucleic acid and protein of steroidogenic enzymes in rat adrenal in vivo 1. *Endocrinology*. 1998;139:3913–22.
6. Raha S, Robinson BH. Mitochondria, oxygen free radicals, disease and ageing. *Trends Biochem Sci*. 2000;25:502–8.
7. Hanukoglu I. Antioxidant protective mechanisms against reactive oxygen species (ROS) generated by mitochondrial P450 systems in steroidogenic cells. *Drug Metab Rev*. 2006;38:171–96.
8. Kil IS, Lee SK, Ryu KW, Woo HA, Hu M-C, Bae SH, Rhee SG. Feedback control of adrenal steroidogenesis via H₂O₂-dependent, reversible inactivation of peroxiredoxin III in mitochondria. *Mol Cell*. 2012;46:584–94.
9. Beattie MC, Chen H, Fan J, Papadopoulos V, Miller P, Zirkin BR. Aging and luteinizing hormone effects on reactive oxygen species production and DNA damage in rat Leydig cells. *Biol Reprod*. 2013;88:100.
10. Abidi P, Leers-Sucheta S, Cortez Y, Han J, Azhar S. Evidence that age-related changes in p38 MAP kinase contribute to the decreased steroid production by the adrenocortical cells from old rats. *Aging Cell*. 2008;7:168–78.
11. Dröge W. Free radicals in the physiological control of cell function. *Physiol Rev*. 2002;82:47–95.
12. Ray PD, Huang B-W, Tsuji Y. Reactive oxygen species (ROS) homeostasis and redox regulation in cellular signaling. *Cell Signal*. 2012;24:981–90.
13. Sena LA, Chandel NS. Physiological roles of mitochondrial reactive oxygen species. *Mol Cell*. 2012;48:158–67.
14. BEHRMAN HR, ATEN RF. Evidence That Hydrogen Peroxide Blocks Hormone-Sensitive Cholesterol Transport into Mitochondria of Rat Luteal Cells*. *Endocrinology*. 1991;128:2958–66.
15. Stocco DM, Wells J, Clark BJ. The effects of hydrogen peroxide on steroidogenesis in mouse Leydig tumor cells. *Endocrinology*. 1993;133:2827–32.
16. Diemer T, Allen JA, Hales KH, Hales DB. Reactive oxygen disrupts mitochondria in MA-10 tumor Leydig cells and inhibits steroidogenic acute regulatory (StAR) protein and steroidogenesis. *Endocrinology*. 2003;144:2882–91.
17. McClung JM, Judge AR, Powers SK, Yan Z. p38 MAPK links oxidative stress to autophagy-related gene expression in cachectic muscle wasting. *Am J Phys Cell Phys*. 2010;298:C542–9.
18. Gutiérrez-Uzquiza Á, Arechederra M, Bragado P, Aguirre-Ghiso JA, Porras A. p38 α Mediates Cell Survival in Response to Oxidative Stress via Induction of Antioxidant Genes EFFECT ON THE p70S6K PATHWAY. *J Biol Chem*. 2012; 287:2632–42.
19. Yamada T, Egashira N, Bando A, Nishime Y, Tonogai Y, Imuta M, Yano T, Oishi R. Activation of p38 MAPK by oxidative stress underlying epirubicin-induced vascular endothelial cell injury. *Free Radic Biol Med*. 2012;52:1285–93.
20. Cuenda A, Rousseau S. p38 MAP-kinases pathway regulation, function and role in human diseases. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Res*. 2007;1773:1358–75.
21. Cuadrado A, Nebreda AR. Mechanisms and functions of p38 MAPK signalling. *Biochem J*. 2010;429:403–17.
22. Zaidi SK, Shen W-J, Bittner S, Bittner A, McLean MP, Han J, Davis RJ, Kraemer FB, Azhar S. p38 MAPK regulates steroidogenesis through transcriptional repression of STAR gene. *J Mol Endocrinol*. 2014;53:1–16.
23. Abidi P, Zhang H, Zaidi SM, Shen W-J, Leers-Sucheta S, Cortez Y, Han J, Azhar S. Oxidative stress-induced inhibition of adrenal steroidogenesis requires participation of p38 mitogen-activated protein kinase signaling pathway. *J Endocrinol*. 2008;198:193–207.
24. Rainey WE, Saner K, Schimmer BP. Adrenocortical cell lines. *Mol Cell Endocrinol*. 2004;228:23–38.
25. Yasumura Y, Buonassisi V, Sato G. Clonal analysis of differentiated function in animal cell cultures. *Cancer research*. 1966;26(3 Part 1):529-35.
26. Temel RE, Trigatti B, DeMattos RB, Azhar S, Krieger M, Williams DL. Scavenger receptor class B, type I (SR-BI) is the major route for the delivery of high density lipoprotein cholesterol to the steroidogenic pathway in cultured mouse adrenocortical cells. *Proc Natl Acad Sci U S A*. 1997;94:13600–5.
27. Hu Z, Li J, Kuang Z, Wang M, Azhar S, Guo Z. Cell-Specific Polymorphism and Hormonal Regulation of DNA Methylation in Scavenger Receptor Class B, Type I. *DNA Cell Biol*. 2016;35:280–9.
28. Reaven E, Tsai L, Azhar S. Intracellular events in the “selective” transport of lipoprotein-derived cholesteryl esters. *J Biol Chem*. 1996;271:16208–17.
29. Hu Z, Hu J, Zhang Z, Shen W-J, Yun CC, Berlot CH, Kraemer FB, Azhar S. Regulation of expression and function of scavenger receptor class B, type I (SR-BI) by Na⁺/H⁺ exchanger regulatory factors (NHERFs). *J Biol Chem*. 2013; 288:11416–35.
30. Zhou Y, Wang Y, Jiaping YU. Research on the determination of serum steroid hormones by isotope dilution HPLC-MS/MS. 2015.
31. Kyriakis JM, Avruch J. Mammalian mitogen-activated protein kinase signal transduction pathways activated by stress and inflammation. *Physiol Rev*. 2001;81:807–69.
32. McCubrey JA, LaHair MM, Franklin RA. Reactive oxygen species-induced activation of the MAP kinase signaling pathways. *Antioxid Redox Signal*. 2006;8:1775–89.
33. Son Y, Kim S, Chung H-T, Pae H-O. Reactive oxygen species in the activation of MAP kinases. *Methods Enzymol*. 2013;528:27–48.
34. Devasagayam T, Tilak J, Boloor K, Sane KS, Ghaskadbi SS, Lele R. Free radicals and antioxidants in human health: current status and future prospects. *Japi*. 2004;52:4.
35. Xing J, Kornhauser JM, Xia Z, Thiele EA, Greenberg ME. Nerve growth factor activates extracellular signal-regulated kinase and p38 mitogen-activated protein kinase pathways to stimulate CREB serine 133 phosphorylation. *Mol Cell Biol*. 1998;18:1946–55.
36. Delghandi MP, Johannessen M, Moens U. The cAMP signalling pathway activates CREB through PKA, p38 and MSK1 in NIH 3 T3 cells. *Cell Signal*. 2005;17:1343–51.
37. Iordanov M, Bender K, Ade T, Schmid W, Sachsenmaier C, Engel K, Gaestel M, Rahmsdorf H, Herrlich P. CREB is activated by UVC through a p38/HOG-1-dependent protein kinase. *EMBO J*. 1997;16:1009–22.
38. Naqvi S, Martin KJ, Arthur JS. CREB phosphorylation at Ser133 regulates transcription via distinct mechanisms downstream of cAMP and MAPK signalling. *Biochem J*. 2014;458:469–79.
39. Quinn P, Payne A. Steroid product-induced, oxygen-mediated damage of microsomal cytochrome P-450 enzymes in Leydig cell cultures. Relationship to desensitization. *J Biol Chem*. 1985;260:2092–9.
40. Grewal T, de Diego I, Kirchoff MF, Tebar F, Heeren J, Rinninger F, Enrich C. High density lipoprotein-induced signaling of the MAPK pathway involves scavenger receptor type BI-mediated activation of Ras. *J Biol Chem*. 2003; 278:16478–81.
41. Pan B, Ma Y, Ren H, He Y, Wang Y, Lv X, Liu D, Ji L, Yu B, Wang Y. Diabetic HDL is dysfunctional in stimulating endothelial cell migration and proliferation due to down regulation of SR-BI expression. *PLoS One*. 2012;7:e48530.
42. Mineo C, Shaul PW. Regulation of signal transduction by HDL. *J Lipid Res*. 2013;54:2315–24.
43. Shaywitz AJ, Greenberg ME. CREB: a stimulus-induced transcription factor activated by a diverse array of extracellular signals. *Annu Rev Biochem*. 1999;68:821–61.
44. Mayr B, Montminy M. Transcriptional regulation by the phosphorylation-dependent factor CREB. *Nat Rev Mol Cell Biol*. 2001;2:599–609.
45. Sakamoto KM, Frank DA. CREB in the pathophysiology of cancer: implications for targeting transcription factors for cancer therapy. *Clin Cancer Res*. 2009;15:2583–7.
46. Stocco DM. StAR protein and the regulation of steroid hormone biosynthesis. *Annu Rev Physiol*. 2001;63:193–213.
47. Simpson ER, Miller DA. Cholesterol side-chain cleavage, cytochrome P450, and iron-sulfur protein in human placental mitochondria. *Arch Biochem Biophys*. 1978;190:800–8.
48. Lavoie HA, King SR. Transcriptional regulation of steroidogenic genes: STAR1, CYP11A1 and HSD3B. *Exp Biol Med*. 2009;234:880–907.
49. Chen H, Irizarry RA, Luo L, Zirkin BR. Leydig cell gene expression: effects of age and caloric restriction. *Exp Gerontol*. 2004;39:31–43.
50. Fu H, Wada-Hiraike O, Hirano M, Kawamura Y, Sakurabashi A, Shirane A, Morita Y, Isono W, Oishi H, Koga K. SIRT3 positively regulates the expression of folliculogenesis-and luteinization-related genes and progesterone secretion by manipulating oxidative stress in human luteinized granulosa cells. *Endocrinology*. 2014;155:3079–87.
51. Indo HP, Yen H-C, Nakanishi I, Matsumoto K-i, Tamura M, Nagano Y, Matsui H, Gusev O, Cornette R, Okuda T. A mitochondrial superoxide theory for oxidative stress diseases and aging. *J Clin Biochem Nutr*. 2015;56:1.

52. Candas D, Li JJ. MnSOD in oxidative stress response-potential regulation via mitochondrial protein influx. *Antioxid Redox Signal*. 2014;20:1599–617.
53. Stocco DM, Wang X, Jo Y, Manna PR. Multiple signaling pathways regulating steroidogenesis and steroidogenic acute regulatory protein expression: more complicated than we thought. *Mol Endocrinol*. 2005;19:2647–59.
54. Lin Y, Hou X, Shen WJ, Hanssen R, Khor VK, Cortez Y, Roseman AN, Azhar S, Kraemer FB. SNARE-mediated cholesterol movement to mitochondria supports steroidogenesis in rodent cells. *Mol Endocrinol*. 2016;30(2):234–47.
55. Vitale G, Salvioli S, Franceschi C. Oxidative stress and the ageing endocrine system. *Nat Rev Endocrinol*. 2013;9:228–40.
56. Azhar S, Cao L, Reaven E. Alteration of the adrenal antioxidant defense system during aging in rats. *J Clin Investig*. 1995;96:1414.
57. Valko M, Leibfritz D, Moncol J, Cronin MT, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol*. 2007;39:44–84.
58. Ye Z-W, Zhang J, Townsend DM, Tew KD. Oxidative stress, redox regulation and diseases of cellular differentiation. *Biochimica et Biophysica Acta (BBA)-General Subjects*. 2015;1850:1607–21.
59. Karlsson H, Kontush A, James RW. Functionality of HDL: antioxidant and detoxifying effects. In *High Density Lipoproteins*. Springer; 2015: 207–228. https://link.springer.com/chapter/10.1007%2F978-3-319-09665-0_5.
60. Kulisz A, Chen N, Chandel NS, Shao Z, Schumacker PT. Mitochondrial ROS initiate phosphorylation of p38 MAP kinase during hypoxia in cardiomyocytes. *Am J Phys Lung Cell Mol Phys*. 2002;282:L1324–9.
61. Yu F-Q, Han C-S, Yang W, Jin X, Hu Z-Y, Liu Y-X. Activation of the p38 MAPK pathway by follicle-stimulating hormone regulates steroidogenesis in granulosa cells differentially. *J Endocrinol*. 2005;186:85–96.
62. Butler MP, Hanly JA, Moynagh PN. Pellino3 is a novel upstream regulator of p38 MAPK and activates CREB in a p38-dependent manner. *J Biol Chem*. 2005;280:27759–68.
63. Jinlian L, Yingbin Z, Chunbo W. p38 MAPK in regulating cellular responses to ultraviolet radiation. *J Biomed Sci*. 2007;14:303–12.
64. Olala LO, Choudhary V, Johnson MH, Bollag WB. Angiotensin II-induced protein kinase D activates the ATF/CREB family of transcription factors and promotes StAR mRNA expression. *Endocrinology*. 2014;155:2524–33.
65. Saha B, Singh SK, Sarkar C, Bera R, Ratha J, Tobin DJ, Bhadra R. Activation of the Mitf promoter by lipid-stimulated activation of p38-stress signalling to CREB. *Pigment Cell Res*. 2006;19:595–605.
66. Manna P, Eubank D, Lalli E, Sassone-Corsi P, Stocco D. Transcriptional regulation of the mouse steroidogenic acute regulatory protein gene by the cAMP response-element binding protein and steroidogenic factor 1. *J Mol Endocrinol*. 2003;30:381–97.
67. Clem BF, Hudson EA, Clark BJ. Cyclic adenosine 3', 5'-monophosphate (cAMP) enhances cAMP-responsive element binding (CREB) protein phosphorylation and phospho-CREB interaction with the mouse steroidogenic acute regulatory protein gene promoter. *Endocrinology*. 2005; 146:1348–56.
68. Cherradi N, Bideau M, Arnaudeau S, Demaurex N, James RW, Azhar S, Capponi AM. Angiotensin II promotes selective uptake of high density lipoprotein cholesterol esters in bovine adrenal glomerulosa and human adrenocortical carcinoma cells through induction of scavenger receptor class B type I. *Endocrinology*. 2001;142:4540–9.
69. Xing Y, Cohen A, Rothblat G, Sankaranarayanan S, Weibel G, Royer L, Francone OL, Rainey WE. Aldosterone Production in Human Adrenocortical Cells Is Stimulated by High-Density Lipoprotein 2 (HDL2) through Increased Expression of Aldosterone Synthase (CYP11B2). *Endocrinology*. 2011;152: 751–63.
70. Saha S, Bornstein SR, Graessler J, Koprassch S. Very-low-density lipoprotein mediates transcriptional regulation of aldosterone synthase in human adrenocortical cells through multiple signaling pathways. *Cell Tissue Res*. 2012;348:71–80.
71. Manna PR, Stocco DM. The Role of Specific Mitogen-Activated Protein Kinase Signaling Cascades in the Regulation of Steroidogenesis. *J Signal Transduction*. 2011;2011:821615.
72. Svechnikov K, Stocco DM, Söder O. Interleukin-1 α stimulates steroidogenic acute regulatory protein expression via p38 MAP kinase in immature rat Leydig cells. *J Mol Endocrinol*. 2003;30:59–67.
73. Tajima K, Dantes A, Yao Z, Sorokina K, Kotsuji F, Seger R, Amsterdam A. Down-regulation of steroidogenic response to gonadotropins in human and rat preovulatory granulosa cells involves mitogen-activated protein kinase activation and modulation of DAX-1 and steroidogenic factor-1. *J Clin Endocrinol Metab*. 2003;88:2288–99.
74. Otis M, Campbell S, Payet MD, Gallo-Payet N. Angiotensin II stimulates protein synthesis and inhibits proliferation in primary cultures of rat adrenal glomerulosa cells. *Endocrinology*. 2005;146:633–42.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

